

# Do Postoperative Drains after Emergency Laparotomy Prevent Deep Surgical Site Infection?

SHABAB HUSSAIN<sup>1</sup>, MUHAMMAD TANVIR IQBAL<sup>2</sup>, ZUBAIR AHMAD<sup>3</sup>, UMAR FAROOQ DAR<sup>4</sup>

## ABSTRACT

**Background:** Emergency laparotomy is a common procedure in our tertiary care hospitals but there is contradictory evidence regarding whether to put postoperative drains or not. Current study was planned to generate the evidence.

**Aim:** To compare the frequency of deep surgical site infection in patients undergoing emergency laparotomy with and without postoperative drains in a tertiary care hospital.

**Methods:** This randomised control trial study was carried out at Surgical Unit of Postgraduate Medical Institute, Lady Reading Hospital, Peshawar over a period of six months. Seven hundred and sixty two patients undergoing emergency laparotomy with age from 15-60 years, due to perforated appendix, tuberculosis, typhoid determined history, clinical examination and erect abdominal X ray were divided into two groups i.e., with and without post-operative drains. Rate of deep surgical site infection on 3<sup>rd</sup> & 7<sup>th</sup> day was measured as outcome.

**Results:** The mean age of the patients was 28.92±6.246 years with 330(43.3%) were female while rest 432(56.7%) were male. Deep surgical site infection on 3<sup>rd</sup> day was 7.2% in patients with post-operative drains while 8.1% in patients without post-operative drains. Similarly on day 7, deep surgical site infection was 6.3% in patients with post-operative drains as compared with 8.1% in patients without post-operative drains. Differences were statistically non-significant.

**Conclusion:** It is concluded that there is no difference in frequency of developing deep surgical site infection on 3<sup>rd</sup> day and 7<sup>th</sup> whether you use post-operative drains after emergency laparotomy or not. So we accept the null hypothesis and conclude the use of post-operative drains is not associated with deep surgical site infection on 3<sup>rd</sup> and 7<sup>th</sup> day.

**Keywords:** Deep surgical site infection, Emergency laparotomy, Postoperative drains, Peritonitis

---

## INTRODUCTION

Prophylactic drainage of the peritoneal cavity after gastro-intestinal (GI) surgery has been used since time immemorial, with the dictum of Lawson Tait, the 19<sup>th</sup> century British surgeon, "when in doubt, drain", well known to all surgical trainees<sup>1,2,3</sup>. Emergency laparotomy is a common procedure in our settings. To drain or not to drain has been a dilemma. Postoperative drains help the surgeon not only to detect anastomosis leakage early but also reduce postoperative adhesions. But on the other hand, drains are associated with deep surgical site infection (DSSI)<sup>4,5</sup>. Deep surgical site infection is among common morbidities ranging from delayed healing to systemic sepsis having high impact on the economy and health care resources<sup>6</sup> due to increased length of stay<sup>7</sup>. In a Pakistani study conducted in tertiary care hospital, the overall rate of surgical site infection came out 13%<sup>8</sup>, much higher than other developed countries like 1.9% in USA<sup>6</sup>.

The available evidence is lacking consensus regarding use of post operative drainage in GI procedures. In a study incidence of DSSI was significantly higher in patients who received a drain (31% vs. 9%, p=0.001).<sup>9</sup> But in another study there came out statistically non-significant difference in the rate of DSSI based on the presence or absence of an intra-abdominal drain after laparotomy (17 vs 18%, P=0.88).<sup>10</sup> Similarly in a third study one drain placement was found as good as the two drain placement.<sup>1</sup> In a retrospective review to determine safety and effectiveness of routine drainage and nondrainage, no significant difference in mean time for return of bowel function (3.8 vs 4.0 days; P=0.6), rate of surgical site infection (63% vs 70%; P=0.39), wound dehiscence (36% vs 27%; P=0.27), anastomotic leak (2.5% vs 1.5%; P=0.27), enterocutaneous fistula formation (10% vs 6.1%; P=0.40), intra-abdominal abscess formation (4% vs 9%; P=0.18), or mean length of hospital stay (22 vs 19 days; P=0.26) was observed<sup>11</sup>.

Deep surgical site infections pose a major threat in all surgical interventions. Abdominal infections are common in our setting because of lack of implementation of standardized protocols for infection control. Gut leakage and post-operative infected secretions lead to deep infections and abscess

---

<sup>1</sup>Dept of Surgery, Lady Reading Hospital Peshawar, <sup>2</sup>Specialist General Surgeon Najran University Hospital Najran KSA, <sup>3</sup>Consultant Surgeon. DHQ Teaching Hospital D Gi Khan, <sup>4</sup>Dept of Infectious diseases, Institute of Public Health Lahore, Correspondence to Dr. Umar Farooq Dar Email: umardar84@gmail.com

formation. Placement of intra-abdominal drains has been a practice in our settings conventionally. The evidence of its benefit is contradictory as explained earlier. Current study aims to explore the better practice regarding placement of drain in term of lower rate of deep surgical site infection. Deep surgical site infection is a common cause of prolonged hospital stay in our already burdened teaching hospitals. Results of this study will help patients achieve health early and hospital managers may get reduction in bed occupancy rate.

**PATIENTS AND METHODS**

This randomised control trial study was carried out at Surgical Unit of Postgraduate Medical Institute, Lady Reading Hospital, Peshawar over a period of six months. Seven hundred and sixty two patients undergoing emergency laparotomy with age from 15-60 years, due to perforated appendix, tuberculosis, typhoid determined history, clinical examination and erect abdominal X ray were divided into two groups i.e. with intra-abdominal drains and without post-operative drains. Patients will be followed post operatively at 3rd day and 7th day for presence of deep surgical site infection by researcher himself. The data was entered and analysed in the SPSS version 17. Chi square test of homogeneity will be applied to determine statistical difference in both groups regarding rate of DSSI on 3<sup>rd</sup> and 7<sup>th</sup> post-operative day. A value of p <0.05 will be considered as significant.

**RESULTS**

There were 330(43.3%) were female while rest 432(56.7%) were male patients. Seven hundred and sixteen (94%) patients were below 40 years while rest of 46 (6%) patients were either 40 or above 40 years of their age with mean age of the patients were 28.92±6.246 ranging from 21 to 59 years. Five hundred and thirty eight 538(70.6%) patients stayed in hospital less than five days while 224(29.4%) patients stayed in hospital five and more than five days. Their hospital stay was between 2 to 9 days with mean of 5.01±1.58 days (Table 1). Among 762 patients, 102(13.4%) patients have deep surgical site infection on 3<sup>rd</sup> day while 55(7.2%) patients showed up with deep surgical site infection on 7<sup>th</sup> day of operation. Deep surgical site infection on 3<sup>rd</sup> day was 7.2% in patients with post-operative drains while 8.1% in patients without post-operative drains. Difference was statistically non-significant. Similarly on day 7, deep surgical site infection was 6.3% in patients with post-operative drains as compared with

8.1% in patients without post-operative drains (Table 2).

Table 1: Demographic information of the patients

Variable	No.	%
<b>Gender</b>		
Male	330	43.3
Female	432	56.7
<b>Age (years)</b>		
<40	716	94.0
≥40	46	6.0
<b>Hospital stay (days)</b>		
<5	538	70.6
≥5	224	29.4

Table 2: Comparison of surgical site infection on 3<sup>rd</sup> and 7<sup>th</sup> of operation

Surgical site infection	Post-operative drains	Without post-operative drains
<b>3<sup>rd</sup> day</b>		
Yes	55 (7.2%)	64 (8.1%)
No	707 (92.8%)	698 (91.9%)
<b>7<sup>th</sup> day</b>		
Yes	47 (6.3%)	64 (8.1%)
No	715 (93.7%)	698 (91.9%)

P>0.05

**DISCUSSION**

Deep surgical site infection (DSSI) is among common morbidities ranging from delayed healing to systemic sepsis having high impact on the economy and health care resources<sup>6</sup> due to increased length of stay.<sup>7</sup> In a Pakistani study conducted in tertiary care hospital, the overall rate of surgical site infection came out 13%<sup>8</sup>, much higher than other developed countries like 1.9% in USA<sup>6</sup>. The available evidence is lacking consensus regarding use of post operative drainage in GI procedures. In a previous study incidence of DSSI was significantly higher in patients who received a drain (31% vs. 9%, p = 0.001)<sup>3</sup>.

In our study, deep surgical site infection on 3<sup>rd</sup> day was 7.2% in patients with post-operative drains while 8.1% in patients without post-operative drains. The difference was statistically non-significant (Table 2). We may conclude that there is no difference in frequency of developing deep surgical site infection on 3<sup>rd</sup> day whether you use post-operative drains after emergency laparotomy or not. Similarly on day 7, deep surgical site infection was 6.3% in patients with post-operative drains as compared with 8.1% in patients without post-operative drains. The difference was statistically non-significant (Table 2). We may conclude that there is no difference in frequency of developing deep surgical site infection on 7<sup>th</sup> day whether you use post-operative drains after emergency laparotomy or not.

## CONCLUSION

It is concluded that there is no difference in frequency of developing deep surgical site infection on 3<sup>rd</sup> day and 7<sup>th</sup> whether you use post-operative drains after emergency laparotomy or not. So we accept the null hypothesis and conclude the use of post-operative drains is not associated with deep surgical site infection on 3<sup>rd</sup> and 7<sup>th</sup> day.

## REFERENCES

1. Ansari MM, Akhtar A, Haleem S, Husain M, Kumar A. Is there a role of abdominal drainage in primarily repaired perforated peptic ulcers? *J Exp Integr Med* 2012; 2(1):47-54.
2. Sharma L, Singh A, Bhaskaran S, Radhika AG, Radhakrishnan G. Fallopian tube herniation: an unusual complication of surgical drain. *Obstet Gynecol* 2012;2012:194350.
3. Mosley JG, Jantet G. Herniation at the site of an abdominal drain. *Br J Clin Prac* 1978; 32(2): 56–8.
4. Malinoski DJ, Patel MS, Yakar DO, Green D, Qureshi F, Inaba K, Brown CV, Salim A. A diagnostic delay of 5 hours increases the risk of death after blunt hollow viscus injury. *J Trauma* 2010;69(1):84-7.
5. Reiffel AJ, Barie PS, Spector JA. A multi-disciplinary review of the potential association between closed-suction drains and surgical site infection. *Surg Infect (Larchmt)* 2013; 14(3): 244-69.
6. Yi M, Edwards JR. Improving risk-adjusted measures of surgical site information for the National Healthcare Safety Network. *Infect Control Hosp Epidemiol* 2011; 2(10):970.
7. Jenks PJ, Laurent M, McQuarry S, Watkins R. Clinical and economic burden of surgical site infection (SSI) and predicted financial consequences of elimination of SSI from an English hospital. *J Hosp Infect* 2013; 85(1): 24-33.
8. Sangrasi AK, Leghari AA, Memon A, Talpur AK, Qureshi GA, Memon JM. Surgical site infection rate and associated risk factors in elective general surgery at a public sector medical university in Pakistan. *Int Wound J* 2008; 5:74–8.
9. Talving P, Mohseni S, Inaba K, Plurad D, Branco BC, Lam L, et al. Closed suction drain after isolated hollow viscus injury: a friend or foe? *J Trauma* 2011; 70(6): 1424-8.
10. Mohseni S, Talving P, Kobayashi L, Kim D, Inaba K, Lam L, et al. Closed-suction drain placement at laparotomy in isolated solid organ injury is not associated with decreased risk of deep surgical site infection. *Am Surg* 2012; 78(10): 1187-91.
11. Nasir AA, Abdur-Rahman LO, Adeniran JO. Is intraabdominal drainage necessary after laparotomy for typhoid intestinal perforation? *J Pediatr Surg* 2012; 47(2): 355-8.