

# Frequency and Determinants of Intestinal Parasitic Infections in Patients Attending Tertiary Care Hospitals

MUHAMMAD ANWAR<sup>1</sup>, MUHAMMAD WAQAR REHAN<sup>2</sup>, WASEEM AHSEN<sup>3</sup>

## ABSTRACT

**Background:** The condition of human in which intestinal tract infected by parasite is called intestinal parasite infection. These parasites can live anywhere in body but prefer the intestinal wall.

**Aim:** To determine the prevalence of gastrointestinal disease patients having IP (intestinal parasitic) infection, associated risk factors and clinical observation..

**Method:** This comparative and observational research was carried out at Ch. Rehmat Ali Trust Hospital during the period from 1<sup>st</sup> January 2017 to 31<sup>st</sup> December 2017. Total 235 patients of both genders with symptoms of acute gastroenteritis were selected. Patients detailed history of sex, age, occupation, literacy level, financial status etc was examine. Symptoms associated with IP infection were also examined. Patients having bloody diarrhea and nosocomial gastroenteritis acquired in hospitalization for other disease were excluded from study.

**Results:** There were 130 (55.32%) patients were male, 105(44.68%) patients were females. 88(37.45%) patients were aged of <24 years, in which 38 patients was having IP infection.102(43.40%) were aged between 25 to 49 years, in which 26 patients found IP infected, and 45 (19.15%) having age above than 49 years, in which 26 patients having IP infections. 135 (57.45%) patients were literate, in which 45% having IP infections and 100 (42.55%) were illiterate, in which 55% having IP infections. IP (Intestinal parasites) was observed in 90 patients, in which, type of Eba\_H (entamoeba histolytica) parasite was mostly found in 30(12.77%). Patients having monthly income ≤25,000 rupees had the highest prevalence of IP infection 41.42%.

**Conclusion:** Significant association of parasitic intestinal infection was found in patients with diarrhea and dysentery in different ages, educational status and monthly income.

**Keywords:** Risk factors, Intestinal, parasitic infection,

---

## INTRODUCTION

According to World Health Organization (WHO), among infants and young children the prevalence of intestinal parasites is high and mostly in those who living in poor communities.<sup>1</sup> Inadequate sanitation and hygiene, improper hygiene, infected soil and consumption of contaminated water are the most common causes of intestinal parasite reported by Ziegelbauer et al.<sup>2</sup> There is lack of facilities in poorest communities to dispose of human waste which cause intestinal parasite infection as well as other diseases. Poor habits of hygiene and non-availability of hygiene resources like facilities of hand washing also impact negatively on the rate of parasite infections<sup>3</sup>.

Parasite infection is reported like a leading cause of non-bacterial gastroenteritis (stomach and intestinal tract infection causing vomiting and diarrhea) particularly among infants and young children globally.<sup>4</sup> Parasite infectivity is associated with infants and younger children increased hospitalizations and deaths<sup>5</sup>.

As per World Health Organization estimation, 215,000 child mortalities took place worldwide during 2013 caused by parasitic infection compared to 528,000 during 2000. Parasite infection national estimates attributable mortalities in under-five children ranged from 47,100 (India) to below 5 mortalities (79 countries). Among all mortalities in India, 22% occurred in children less than 5 years old. Four countries (Pakistan, India, Congo and Nigeria) accounted

virtually half (49 percent) of all mortalities under-five during 2013<sup>6</sup>. Prevalence of intestinal parasitic infection in Pakistan is estimated between 20 to 40 percent<sup>7</sup>.

Protozoans and helminthes are the major groups of parasites. Of these protozoans including isospora, microsporidia and cryptosporidium are much common in HIV infected. These parasites can infect the intestinal tract. Among developing countries, parasite is most common recognized pathogen among children with acute gastroenteritis less than 2 years of age. Among developed states, parasites have been indentified among 35 to 50 percent infants and young hospitalized children with severe diarrhea<sup>8</sup>.

The primary transmission mode is through fecal-oral route with symptoms normally developing after an incubation phase of one to two days. Most of the children got infected with parasite during their first 3 years of life, with maximum prevalence of diarrhea between six to twenty four months. Prior infectivity gives safeguard from later illnesses. Although the re-infections are common but later illnesses tend to be less acute<sup>9</sup>. Diarrhea attributable to parasites could be caused by impaired glucose and sodium assimilation, as harmed cells on villi are restored by non-assimilated undeveloped crypt cells.<sup>10</sup> Better sanitation and hygiene can significantly lower the rate of diarrhea, while diarrhea is just slightly influenced through sanitation and hygiene improvements. It is expected that through vaccination, mortality and morbidity rates of diarrhea due to parasitic infectivity can be decreased among children. Among countries that have conducted a program regarding vaccination, reductions in the rate of acute parasitic

---

<sup>1</sup>Assistant Professor, Pediatric Medicine, Continental Medical College Lahore,

<sup>2</sup>HO, Medicine, Shalamar Hospital Lahore, <sup>3</sup>Associate Professor of Medicine, Islam Medical College, Sialkot

Correspondence to Dr. Wasim Ahsen Email: drwasemalik@yahoo.com

infectivity have been seen, however mild infectivity remains ongoing.<sup>11</sup>

**METHODS**

It was cross-sectional, observational study conducted at Ch. Rehmat Ali Trust Hospital during the period from 1st January 2017 to 31st December 2017. Total 260 patients both males and females with symptoms of acute gastroenteritis were included in the study. Patients with the bloody diarrhea and nosocomial gastroenteritis acquired in hospitalization for other disease were excluded. Samples of stool were taken immediately within 24-48 hours to prevent nosocomial infection. Containers of stool were labeled with unique patient identifier. Identification of parasitic infections in patient's stool was carried out through enzyme linked immunoassay. Data was analyzed using SPSS version 20. Frequencies and percentages were calculated for all categorical variables. Mean and standard deviation were calculated for age.

**RESULTS**

Out of 235 patients, 130 (55.32%) patients were male, 105 (44.68%) patients were females. In males patients we found 53(40.77%) had IP (intestinal parasitic) infection and out of 105 female patients, we found 37 (35.24%) had IP infection. 88 (37.45%) patients were aged of <24 years, in which 50(56.82%) patients had IP infection. 102 (43.40%) patients were aged between 25 to 49 years, in which 22 (21.57%) patients found IP infected, and 45 (19.15%) patients having age above than 49 years, in which 18 (40%) patients had IP infections. Intestinal parasites was observed in 90 (38.29%) patients, we found types of IP (intestinal parasites) were entamoeba histolytica (Eba\_H), entamoeba coli (Eba-Coli), giardia lamblia (GI), ascaris ricoide, lba-butchi, hymenolipis nana, taenia species, blastocystis hominis, trichuritic hurra as prevalence of 33 (14.04%), 4 (1.7%), 24 (10.21%), 2(0.85%), 1(0.42%), 19(8.09%), 1(0.42%), 5 (2.13%) and 1 (0.42%) respectively (Table 1).

135 (57.45%) patients were literate and 40(29.63%) having IP infections and 100 (42.55%) were illiterate and 50% had IP infection. 140 (59.57%) patients having monthly income ≤25,000 rupees had the highest prevalence of IP infection 41.42% than the 95 (40.43%) patients having income of > 25000/month with prevalence of IP infection 33.68% in 32 patients. Another common risk factor, use of not-filtered water was observed in 125 patients with frequency of IP infection was 40% and those patients whom were using filtered water the prevalence was noted as 36.36%. We found 130 patients with 43.85% frequency of IP infection was resulted as not washing hands before eating and washing hand before eating found 42.43% prevalence of IP Infection (Table 2). The symptoms associated to IP infection were also examined such as diarrhea, dysentery, stomach pain, Abdominal pain, nausea and vomiting, bloating and constipation as 41.67%, 42.42%, 42.5%, 38%, 33.33%, 35%, and 17.64% respectively (Table 1).

Table 1: Frequency of IP among patients with gastrointestinal disorders (n=260)

Parasites	Frequency No.	%age
Ent-ba histolytica	33	14.04
Ent-ba coli	4	1.70
GI (giardia lamblia)	24	10.21
Ascaris ricoide	2	0.85
lo-ba butchii	1	0.42
Hymenolepis nana	19	8.09
Taenia specie	1	0.42
Blastocystis hominis	5	2.13
Trichhuristic hurra	1	0.42
No IP	145	67.7

Table 2: Risk factors associated with IP among patients with gastrointestinal disorders

Characteristics	F/No.	Found	Not Found	P value
<b>Age (years)</b>				
<24	88	50 (56.82%)	38 (43.18%)	0.01
25-49	102	22 (21.57%)	81 (78.43%)	
>49	45	18 (40.0%)	27 (60.0%)	
<b>Gender</b>				
Male	130	53(40.8%)	77 (59.2%)	0.16
Female	105	37 (35.2%)	68 (64.8%)	
<b>Literacy Level</b>				
Literate	135	40 (29.63%)	95 (70.37%)	0.02
Illiterate	100	50 (50%)	50 (50%)	
<b>Monthly income (Rs.)</b>				
< 25,000	140	58 (41.42%)	82 (58.58%)	0.00
>25,000	95	32 (33.68%)	63 (66.32%)	
<b>Drinking water</b>				
Filtered	110	40 (36.36%)	70 (63.64%)	0.58
Not-filtered	125	50 (40%)	75 (60%)	
<b>Washing hand before eating</b>				
Yes	105	33 (42.43%)	72 (68.57%)	0.47
No	130	57 (43.85%)	73 (56.15%)	

Table 3: Symptoms associated with intestinal parasites among patients with gastrointestinal disorders

Characteristics	No.	Found	Not Found
Abdominal Pain	50	19 (38%)	31 (62%)
Nausea Vomit	15	5 (33.33%)	10 (66.67%)
Stomach Pain	40	17 (42.5%)	23 (57.5%)
Bloating	20	7 (35%)	13 (65%)
Diarrhoea	60	25(41.67%)	35 (58.33%)
Dysentery	33	14(42.42%)	19 (57.58%)
Constipation	17	3 (17.64)	14 (82.36%)

**DISCUSSION**

Globally, parasites infection is more common and a major health issue. To determine the prevalence of gastrointestinal disease patients having IP (intestinal parasitic) infection, determine the associated risk factors and clinical observation regarding IP infection. In this study total 235 patients of both genders with symptoms of acute gastroenteritis were selected. This research showed that prevalence of IP (intestinal parasite) was 38.3% after complete examination of infected patients. Our reseach findings are better than the research conducted by Siddiqui et al<sup>7</sup> at Karachi, their results showed that prevalence of IP was 68.8%. The most frequent parasites were Entamoeba

histolytica, Giardia lamblia and Hymenolepis nana in our study however, if we go through the other study Siddiqui et al resulted that most common IP (intestinal parasites) were Giardia lamblia, Entamoeba histolytica and Ascaris lumbricoides.<sup>5</sup> Another study regarding IP infections was in Lahore by Ghani et al<sup>9</sup> shows better results that IP prevalence was only 21.0% and Enterobius vermicularis, Giardia lamblia and Entamoeba histolytica were the most common parasites.

In This study, we observed that most effected group was more than <24 years old and mostly men were affected. But these results were different from the other study that was conducted by Meraj et al<sup>12</sup> resulted that mostly patients regarding IP infection were women.

Education is most important factor and plays an important role in preventing people from multiple health problems. Our study shows that the prevalence of IP infection in illiterate patients were high than literate patients, these results shows similarity to the other studies Kiani et al<sup>13</sup> also confirmed that parasite infection was most frequent among illiterate patients.

The importance of pure drinking water is the most common factor plays an important role for preventing people from many health related disorders. Our research resulted that IP (intestinal parasite) infection was high among patients who used unhealthy/not filtered water. The findings of our study are comparable with a recent study conducted by dos Banhos and fellows<sup>14</sup> who also confirmed that intestinal parasite infection was more prevalent among patients who used untreated water.

We found 130 patients with 43.85% frequency of IP infection was resulted as not washing hands before eating and washing hand before eating found 42.43% prevalence of IP Infection and this shows similarity to the some other research conducted to evaluate the prevalence of IP infection in patients with gastrointestinal disorders.<sup>14-15</sup>

The symptoms associated to IP infection were also examined such as diarrhea, dysentery, stomach pain as 41.67%, 42.42%, 42.5% respectively, approximately similarity was observed in some other studies.

A short research of study was that our patients was small in numbers, We have to do more work about this research to provide the better treatment to the patients whom had IP infection.

## CONCLUSION

It is concluded that mostly patients were aged < 24 years had IP infection and the prevalence of IP infection was high as compared to the developed countries. The most prevalent parasites were Entamoeba histolytica, Giardia lamblia and Hymenolepis nana. Significant association was found in age, education and monthly income with significant association with diarrhea and dysentery.

Availability of safe drinking water and health department intervention can prevent population from such infections.

## REFERENCES

1. WHO. Intestinal worms. Geneva: World Health Organization. Retrieved 2017-03-28.
2. Ziegelbauer K, Speich B, Mäusezahl D, Bos R, Keiser J, Utzinger J. Effect of sanitation on soil-transmitted helminth infection: systematic review and meta-analysis. *PLoS Med* 2012;9(1):e1001162.
3. Ashtiani MT, Monajemzadeh M, Saghi B, Shams S, Mortazavi SH, Khaki S, Mohseni N, et al Prevalence of intestinal parasites among children referred to Children's Medical Center during 18 years (1991-2008), Tehran, Iran. *Ann Trop Med Parasitol* 2011;105(7):507-12.
4. Khan W, Noor-un-Nisa, Khan A. Prevalence and risk factors associated with intestinal parasitic infections among food handlers of Swat, Khyber Pakhtunkhwa, Pakistan. *J Food Nutr Res* 2017; 5(5): 331-6.
5. Hailegebriel T. Prevalence of intestinal parasitic infections and associated risk factors among students at Dona Berber primary school, Bahir Dar, Ethiopia. *BMC Infect Dis* 2017; 17: 362.
6. Babakhani M, Safari R, Rajati F, Salimi S, Omidandoost A. Prevalence and risk factors associated with intestinal parasitic infections among school children in Gashky, West of Iran. *Int J Pediatr* 2017; 5(7): 5263-73.
7. Siddiqui MI, Bilqees FM, Ilyas M, Perveen S. Prevalence of parasitic infections in a rural area of Karachi, Pakistan. *JPMA* 2002; 52: 31-5.
8. Mumtaz S, Siddiqui H, Ashfaq T. Frequency and risk factors for intestinal parasitic infection in children under five years age at a tertiary care hospital in Karachi. *JPMA* 2009; 59: 216.
9. Ghani JJ, Ahmed N, Ashraf K, Ijaz M, Maqbool A. Prevalence of intestinal parasites from fingernails of primary school going children of district Lahore. *J Parasitol Vector Biol* 2016; 8(12): 122-5.
10. Haque R. Human intestinal parasites. *J Health Popul Nutr* 2007; 25(4): 387-91.
11. World Health Organization (2017). Soil-transmitted helminth infections. Available at: <http://www.who.int/mediacentre/factsheets/fs366/en/>
12. Mehraj V, Hatcher J, Akhtar S, Rafique G, Beg MA. Prevalence and factors associated with intestinal parasitic infection among children in an urban slum of Karachi. *PLoS ONE* 2008; 3(11): e3680.
13. Kiani H, Haghighi A, Salehi R, Azargashb E. Distribution and risk factors associated with intestinal parasite infections among children with gastrointestinal disorders. *Gastroenterol Hepatol Bed Bench* 2016; 9(1): S80-7.
14. dos Banhos EF, da Rocha JAM, Pimentel ML, Batista ETM, Silva LM. Prevalence and risk factors for intestinal parasite infections in schoolchildren, in the city of Santarém, Pará State, Brazil. *ABCS Health Sci* 2017; 42(3): 137-42.
15. Kiani H, Haghighi A, Rostami A, Azargashb E, Seyyed Tabaei SJ, Solgi A, Zebardast N. Prevalence, risk factors and symptoms associated to intestinal parasite infections among patients with gastrointestinal disorders in Nahavand, Western Iran. *Rev Inst Med Trop Sao Paulo* 2016; 58: 42.