

Comparison of Laminoplasty with Laminectomy and fusion in the Management of Multilevel Cervical Spondylotic Myelopathy

MUHAMMAD ADNAN^{*A}, MUHAMMAD JAHANGIR KHAN^A, ATTIQUE UR REHMAN^A, SHAHZAD SHAMS^B, HASSAAN ZAHID^C, SAMI MUMTAZ^A

^aGujranwala Medical College/Teaching Hospital, Gujranwala, Pakistan

^bKing Edwards Medical University/ Mayo Hospital, Lahore, Pakistan

^cChildren's hospital and Institute of Child health, Lahore, Pakistan

Correspondence to Dr. Muhammad Adnan Email: dradnansgr@gmail.com Cell:+923466221857

ABSTRACT

Aim: To compare the functional outcome (improvement in pain and gait) and complication rates in the patients who underwent laminoplasty or laminectomy plus fusion for multilevel cervical spondylotic myelopathy in our hospital. Secondly, the study will also provide comparative analysis of pre-operative statistics including age, gender, and duration and severity of myelopathic symptoms in the studied population.

Methods: The patients who underwent laminoplasty and laminectomy plus fusion for multilevel cervical spondylotic myelopathy at the Department of Neurosurgery, DHQ Teaching hospital, Gujranwala from June 2015 to May 2019 were retrospectively analysed. Information collected included patient's age, gender, duration of symptoms, severity of myelopathy (Nurick grade), operation time, blood lost during surgery, hospital stay duration, objective improvement in patient function/Nurick grade, subjective improvement in pain and gait, and post-operative complications. Statistical analysis was done using SPSS version 25. Independent sample T test and Chi-square test for independence were used for quantitative and qualitative variables respectively to determine their significant association with type of surgery. The p values were statistically significant if < 0.05 .

Results: Out of 48 patients, 28(58.3%) underwent laminectomy with fusion while 20(32.7%) underwent laminoplasty. Both cohorts had similar preoperative data including age ($p=0.884$), gender ($p=0.635$) duration of symptoms ($p=0.918$), and Nurick grade ($p=0.827$). In comparison to laminectomy with fusion group, the patients who underwent laminoplasty had significantly lesser operation time (1.65 ± 0.54 hours vs 2.64 ± 0.54 hours, $p<0.01$), lesser blood loss (198.00 ± 82.44 ml vs 352.68 ± 118.09 ml hours, $p<0.01$), and lesser hospital stay time (3.50 ± 0.83 days vs 4.86 ± 1.27 days, $p<0.01$).

Conclusion: Laminoplasty had lesser operation time, blood loss, and hospital stay time than laminectomy with fusion among patients with multilevel cervical spondylotic myelopathy, however subjective and objective outcomes and complication rates were nearly similar with both procedures, suggesting non-superiority of one over other. Prospective trials with large sample size are required to find whether one procedure is truly superior

Keywords: Cervical spondylotic myelopathy, Laminectomy with fusion, Laminoplasty, SPSS

INTRODUCTION

Cervical spondylotic myelopathy is common in elderly people world-wide¹. It affects upto 5% of people of age more than 40 years². It often involves multiple vertebral levels and results progressive neurologic deterioration³. Now a days, laminectomy with posterior fusion⁴ and laminoplasty⁵ are two surgical modalities utilized for the management of multilevel cervical spondylotic myelopathy. Addition of posterior spinal fusion to laminectomy is essential to escape the patients from the risk of postlaminectomy kyphosis⁶. Both current modalities are not without morbidities. Laminectomy with fusion is associated with increased risk of adjacent segment degeneration. Laminoplasty evolved in Japan in 1970⁷. It results less alteration in spine natural biomechanics, however it is associated with mechanical neck pain. It is also absolutely contraindicated in kyphotic cervical alignment⁸. Our both surgical modalities are associated with some risks and benefits, non is gold standard. Internationally, several studies and meta-analyses are available favouring benefits of both laminectomy plus fusion and laminoplasty in patients with multilevel cervical spondylotic myelopathy,

however superiority of one surgical modality over the other is a subject of controversy in the available literature^{9,10,11}. Secondly, these majority literature findings were from Western population, local studies about comparison of laminectomy and fusion with laminoplasty are scarce. Therefore, the objective of this study was to compare the functional outcome (improvement in pain and gait) and complication rates in the patients who underwent laminoplasty and laminectomy plus fusion for multilevel cervical spondylotic myelopathy in our hospital. Secondly, the study will also provide comparative analysis of pre-operative statistics including age, gender, and duration and severity of myelopathic symptoms in the studied population.

MATERIAL AND METHODS

The patients who underwent laminoplasty or laminectomy plus fusion for multilevel cervical spondylotic myelopathy at the Department of Neurosurgery, DHQ Teaching hospital, Gujranwala from June 2015 to May 2019 were retrospectively analysed. Inclusion criteria were (1) patients with progressive myelopathy or myeloradiculopathy symptoms and signs who failed nonsurgical therapies, (2) multilevel disease with cord compression at ≥ 3 levels on imaging; and (3) lordotic sagittal alignment with absent

Received on 17-10-2019

Accepted on 24-01-2020

large anteriorly located osteophyte. The patients with metabolic disorders, cervical fractures, tumor, instability of cervical segment, history of cervical spine operation, or those with combined laminectomy and laminoplasty were excluded. Information collected included patient's age, gender, duration and severity of myelopathic feature (Nurick grade), operation time, blood lost during surgery, hospital stay duration, objective improvement in patient function/Nurick grade, subjective improvement in pain and gait, and post-operative complications. According to Nurick clinical grading of myelopathy¹², root disease without spinal cord involvement was grade zero, while cord disease without walking difficulty was grade 1. Patients with slight walking difficulty which does not restrict full-time employment were in grade 2, while patients with extreme walking difficulty which restrict full-time employment were in grade 3. The walking difficulty which requires assistance of someone, or walker is grade 4, while bed-ridden or chair bound patient is in grade 5. Each procedure was personalized to the patient's discrete pathology. At least 6 months after surgery, gait and pain improvement was assessed on a on a 5-point scale, where very good, good, fair, poor, or very poor was noted on a descending scale from 1 to 5. A response in area of very good (1) or good (2) for each gait and pain was considered a positive response to surgery. In addition to this subjective measurement, we also noted objective improvement in patient function by measuring improvement in the Nurick grade at 6 months after surgery. Postoperative imagings obtained at 1- and 6-months' follow-up were analyzed for problems with hardware (loose or broken screws), cervical lordosis, and lucent portion around the work area. In addition to that other postoperative complications were also noted. All findings were recorded in a structured performa. Statistical Package for Social Science (SPSS), version 25 was used. Means with standard deviations were computed of quantitative variables, and frequencies-percentages for qualitative variables. Chi-square test for independence and Independent sample T test were used for qualitative and quantitative variables respectively to determine their significant association with type of surgery. The p values were taken statistically significant if < 0.05.

RESULTS

Out of 48 patients with multilevel cervical spondylotic myelopathy, 28 patients underwent laminectomy with fusion while 20 underwent laminoplasty. Both cohorts had similar preoperative data. In both procedure groups, there

was no statistically significant difference of mean age of the patients (64.21±5.79 years vs 63.95±6.59 years, p=0.884), mean duration of symptoms (14.04± 4.43 months vs 13.90± 4.54 months, p=0.918), and mean preoperative Nurick grade (3.36±0.83 vs 3.30±0.92, p=0.827) (Table 1). Similarly, the distribution of gender in both cohorts was similar (p=0.635) (Graph 1, Table 2). In comparison to laminectomy with fusion group, the patients who underwent laminoplasty had significantly lesser operation time (1.65±0.54 hours vs 2.64± 0.54 hours, p<0.01), lesser blood loss (198.00±82.44 ml vs 352.68±118.09ml hours, p<0.01), and lesser hospital stay time (3.50 ± 0.83 days vs 4.86±1.27days, p<0.01) (Table 1). There was a similar result of objective improvement in Nurick grade after laminectomy with fusion and laminoplasty (67.9% vs 65%, p=0.539), and subjective improvement in pain (71.4% vs 70%, p=0.582) and gait (60.7% vs 60%, p=0.597). The incidence of post-surgery complications was also similar in both cohorts (10.7% vs 10%, p=0.660) (Table 2). Chronic pain, instrumentation failure, deep infection, and development of kyphotic alignment were the post-operative complications in laminectomy with fusion group, while chronic pain and persistent radiculopathy were seen in laminoplasty group. One patient in each cohort required revision surgery.

Graph 1: Gender wise distribution of patients who underwent laminectomy fusion and laminoplasty (n=48)




Table 1: Associations of various quantitative variables with type of procedure (Laminectomy with fusion vs Laminoplasty) performed in patients suffering multilevel cervical spondylotic myelopathy (n = 48)^{*}

Quantitative variables	Type of procedure		Mean difference	p-value
	Laminectomy fusion (mean + SD)	Laminoplasty (mean + SD)		
Age (years)	64.21 ± 5.79	63.95 ± 6.59	0.26	0.884
Duration of symptoms (months)	14.04 ± 4.43	13.90 ± 4.54	0.14	0.918
Severity of myelopathy (Nurick grade)	3.36 ± 0.83	3.30 ± 0.92	0.06	0.823
Operation time (hours)	2.64 ± 0.54	1.65 ± 0.54	0.99	<0.01
Blood loss (ml)	352.68 ± 118.09	198.00 ± 82.44	154.168	<0.01
Hospital stay (Days)	4.86 ± 1.27	3.50 ± 0.83	1.36	<0.01

^{*}Independent sample T-test was used

Table 2: Associations of various qualitative variables with type of procedure (Laminectomy with fusion vs Laminoplasty) performed in patients suffering multilevel cervical spondylotic myelopathy (n = 48)

Predictors / Factors	Type of procedure		Total	P value
	Laminectomy fusion	Laminoplasty		
Male	21 (75%)	15 (75%)	36 (75%)	0.635
Female	07 (25%)	05 (25%)	12 (25%)	
Objective improvement in patient function (Nurick grade):				0.539
Yes	19 (67.9%)	13 (65%)	32 (66.7%)	
No	09 (32.1%)	07 (35%)	16 (33.3%)	
Subjective improvement in pain:				0.582
Yes	20 (71.4%)	14 (70%)	34 (70.8%)	
No	08 (28.6%)	06 (30%)	14 (29.2%)	
Subjective improvement in gait:				0.597
Yes	17 (60.7%)	12 (60%)	29 (60.4%)	
No	11 (34.3%)	08 (40%)	19 (39.6%)	
Post-operative complications:				0.660
Yes	03 (10.7%)	02 (10%)	05 (10.4%)	
No	25 (89.3%)	18 (90%)	43 (89.6%)	

Chi-square test for independence was used

DISCUSSION

Cervical spondylotic myelopathy mainly affects male in their 7th decade of life¹³. Michael G. Fehlings et al studied multilevel cervical spondylotic myelopathy patients where dominant gender was male (71.2%) in studied population¹⁴. A study from Egypt also showed that patients suffering from multilevel cervical spondylotic myelopathy who came for surgical decompression belonged to male gender mainly (65.8%).¹⁵ In our study, 75% of the total of cervical spondylotic myelopathy patients were male. Huairong Ding et al¹⁶ studied multilevel cervical spondylotic myelopathy patients who were offered laminoplasty and laminectomy with fusion at Tianjin University hospital, China. Mean age of the patients with multilevel cervical spondylotic myelopathy was 61.9±10.6 years, while gender was dominantly male (76.7%, 89 out of 116). Darryl Lau et al¹⁷ found that in comparison to laminoplasty, laminectomy plus fusion exhibited a higher long-term complication rate (p = 0.036), more perioperative blood loss (p < 0.001), and longer hospital stay (p = 0.054). Mean age of all patients was 63.0 years. Similarly in our study, mean age of the patients was 64.10± 6.068 years which suggest that the disease prevalence in 7th decade globally. In our study, both surgical modalities had comparable clinical outcome findings as well as complication rates. Objective improvement in patients function (a fall in Nurick grade) laminectomy plus fusion and laminoplasty was seen in 67.9% and 65% patients respectively. The association was statistically insignificant (p=0.635). Similarly, subjective improvement in pain occurred in 71.4% and 70% patients respectively after two therapeutic surgical modalities. Subjective improvement in gait also occurred in 60.7%, and 60% patients respectively. Similarly, Chang-Hyun Lee and colleagues observed a similar loss of lordosis as well as clinical improvement in both groups of multilevel cervical spondylotic myelopathy patients managed with laminoplasty and laminectomy with fusion.¹⁸ In a systematic review of 15 studies, insignificant difference was found in the incidence of surgical and neurological complications in multilevel cervical spondylotic myelopathy patients treated with both laminoplasty and laminectomy with fusion.¹⁹ On the other hand, Chiazor U Onyia & Sajesh K Menon demonstrated that laminoplasty have a better

functional outcome compared to laminectomy and fusion²⁰. In a meta-analysis of 23 studies about patients treated with laminectomy followed by fusion and laminoplasty, it was concluded that there was no statistically significant difference in JOA scores improvement rate (p=0.27), pain improvement rate (p=0.94), total complications rates (p=0.07), and blood loss (p=0.51)²¹. A meta-analysis of 14 studies²² revealed no significant difference in terms of postoperative neck pain relief (P=.64), improvement in Nurick grade (P=.16) and reoperation rate (P=.21) between two surgical modalities groups. Compared with laminoplasty group, complication rate (P< .00001) was higher in the laminectomy-fusion group. In our study, patients undergoing laminoplasty had significantly lesser operation time duration (p<0.01), lesser blood loss (p<0.01), and shorter hospital stay duration (p<0.01) than patients undergoing laminectomy with fusion. Similarly, Xiang Lin and his team concluded that laminoplasty had a shorter operation time (P<.05), lesser blood loss (P<.05), and a lower complication rate (P<.05) than laminectomy with fusion²³. In another study, multi level cervical spondylotic myelopathy patients treated with both laminoplasty and laminectomy with fusion achieved good but a statistically similar clinical improvement. However, laminoplasty group showed lower mean operation time (p< 0.001) and mean blood loss (p=0.03) than laminectomy fusion group.²⁴ Hence, both laminoplasty and laminectomy with fusion have excellent clinical improvement and comparable complications and risk. Multicenteric studies with larger sample size may nominate single best modality as the standard of care in patients with multilevel cervical spondylotic myelopathy.

CONCLUSION

Laminoplasty had lesser operation time, blood loss, and hospital stay duration than laminectomy with fusion among patients with multilevel cervical spondylotic myelopathy, however subjective and objective outcomes and complication rates were nearly similar with both procedures, suggesting non-superiority of one over other. Prospective trials with large sample size are required to find whether one procedure is truly superior.

REFERENCES

1. de Oliveira Vilaça C, Orsini M, Leite MA, et al. Cervical Spondylotic Myelopathy: What the Neurologist Should Know. *Neurol Int.* 2016;8(4):6330.
2. Editorial. A focus on patient outcomes in cervical myelopathy. *The Lancet Neurology* 2019; 18(7):615. DOI:[https://doi.org/10.1016/S1474-4422\(19\)30168-1](https://doi.org/10.1016/S1474-4422(19)30168-1).
3. Aljuboori Z, Boakye M. The Natural History of Cervical Spondylotic Myelopathy and Ossification of the Posterior Longitudinal Ligament: A Review Article. *Cureus* 2019;11(7):e5074.
4. Bou Monsef JN, Siemionow KB. Multilevel cervical laminectomy and fusion with posterior cervical cages. *J Craniovertebr Junction Spine.* 2017;8(4):316–321.
5. Cho SK, Kim JS, Overley SC, and Merrill RK. Cervical Laminoplasty: Indications, Surgical Considerations, and Clinical Outcomes. *J Am Acad Orthop Surg.* 2018;26(7): e142-e152.
6. Srivastava S, Marathe N, Bhosale S, and Gautham S. Correction of postlaminectomy cervical kyphosis in an operated case of cervical spine primitive neuroectodermal tumor. *Asian J Neurosurg* 2019;14:1017-20.
7. Hirano Y, Ohara Y, Mizuno J, Itoh Y. History and Evolution of Laminoplasty. *Neurosurg Clin N Am.* 2018;29(1):107-113.
8. <https://emedicine.medscape.com/article/1890493-overview#showall>
9. Jung JM, Chung CK, Kim CH, & Yang SH. Clinical and radiological outcomes of C3–C6 laminoplasty with C7 dome-like laminectomy. *Interdisciplinary Neurosurgery: Advanced Techniques and Case Management* 2019; 15:47-52.
10. Singhatanadgige W, Limthongku W, Valone F, Yingsakmongko W, and Riew KD. Outcomes following Laminoplasty or Laminectomy and Fusion in Patients with Myelopathy Caused by Ossification of the Posterior Longitudinal Ligament: A Systematic Review. *Global Spine J* 2016; 6: 702–709.
11. Nurboja B, Kachramanoglou C, and Choi D. Cervical Laminectomy vs Laminoplasty: Is There a Difference in Outcome and Postoperative Pain? *Neurosurgery* 2012; 70(4):965–970.
12. Ramesh VG, Kannan MG, Sriram K, Balasubramanian C. Prognostication in cervical spondylotic myelopathy: Proposal for a new simple practical scoring system. *Asian J Neurosurg* 2017;12:525-8.
13. Northover JR, Wild JB, Braybrooke J, Blanco J. The epidemiology of cervical spondylotic myelopathy. *Skeletal Radiol.* 2012;41(12):1543-6.
14. Fehlings MG, Kopjar B, Defino HLA, Barbagallo G, Arnold PM, Zileli M et al. Laminoplasty versus Laminectomy and Fusion to Treat Cervical Spondylotic Myelopathy: Outcomes of the Prospective Multicenter AOSpine International CSM Study. *The Spine Journal* 2013; 13(9): S2 - S3
15. Abdalla ANM, Sameer MW, Nageeb AM, El-Shehaby, El-Werdany MA, and Abdalla MAM. Comparative Study between Cervical Laminectomy and Cervical Laminectomy with Fixation on Clinical Outcome and Spine Stability in Patients with Degenerative Cervical Myelopathy. *The Egyptian Journal of Hospital Medicine* 2018;72(6):4620-4624.
16. Ding H, Xue Y, Tang Y, He D, Li Z, Zhao Y, et al. Laminoplasty and Laminectomy Hybrid Decompression for the Treatment of Cervical Spondylotic Myelopathy with Hypertrophic Ligamentum Flavum: A Retrospective Study. *PLoS ONE* 2014;9(4): e95482.
17. Lau D, Winkler EA, Than KD, Chou D, and Mummaneni PV. Laminoplasty versus laminectomy with posterior spinal fusion for multilevel cervical spondylotic myelopathy: influence of cervical alignment on outcomes. *J Neurosurg Spine.* 2017;27(5):508-517.
18. Lee CH, Lee J, Kang JD, Hyun SJ, Kim KJ, Jahng TA, et al. Laminoplasty versus laminectomy and fusion for multilevel cervical myelopathy: a meta-analysis of clinical and radiological outcomes. *J Neurosurg Spine* 2015; 22 (6):589–595,
19. Lao L, Zhong G, Li X, Qian L, Liu Z. Laminoplasty versus laminectomy for multi-level cervical spondylotic myelopathy: a systematic review of the literature. *J Orthop Surg Res.* 2013;8:45.
20. Onyia CU, Menon SK. Laminectomy versus laminoplasty in the surgical management of long-segment intradural spinal tumors: Any difference in neurological outcomes? *Asian J Neurosurg* 2018;13:1128-33.
21. Liu FY, Yang SD, Huo LS, Wang T, Yang DL, Ding WY. Laminoplasty versus laminectomy and fusion for multilevel cervical compressive myelopathy: A meta-analysis. *Medicine (Baltimore).* 2016;95(23): e3588.
22. Yuan X, Wei C, Xu W, Gan X, Cao S, Luo J. Comparison of laminectomy and fusion vs laminoplasty in the treatment of multilevel cervical spondylotic myelopathy: A meta-analysis. *Medicine (Baltimore).* 2019;98(13): e14971.
23. Lin X, Cai J, Qin C, Yang Q, and Xiao Z. Comparison of clinical outcomes and safety between laminectomy with instrumented fusion versus laminoplasty for the treatment of multilevel cervical spondylotic myelopathy. *Medicine (Baltimore).* 2019;98(8): e14651.
24. Ma L, Liu FY, Huo LS, Zheng-Qi Z, Xian-Ze S, Feng L, et al. Comparison of laminoplasty versus laminectomy and fusion in the treatment of multilevel cervical ossification of the posterior longitudinal ligament: A systematic review and meta-analysis. *Medicine* 2018; 97(29): e11542.