

Comparison of Titanium cage with PEEK Cage in anterior Cervical Discectomy and fusion (ACDF) Surgery

MUHAMMAD ADNAN^A, MUHAMMAD JAHANGIR KHAN^A, ATTIQUE UR REHMAN^A, HASSAAN ZAHID^C, SHAHZAD SHAMS^B, SAMINA ASGHAR

^a Gujranwala Medical College/Teaching Hospital, Gujranwala, Pakistan

^b King Edwards Medical University/ Mayo Hospital, Lahore, Pakistan

^c Children's hospital and Institute of Child health, Lahore, Pakistan

Correspondence to Dr. Muhammad Adnan

ABSTRACT

Aim: To compare the Titanium cage with PEEK cage in patients who underwent anterior cervical discectomy and fusion (ACDF) surgery at neurosurgery department of DHQ hospital, Gujranwala.

Methods: This was a retrospective analysis of the patients who underwent anterior cervical discectomy and fusion (ACDF) surgery from January 2015 to December 2018 at DHQ Teaching hospital, Gujranwala. The patients among which titanium cage was used were labelled as group A, while those in which PEEK cage was used, were labelled as group B. Information composed from both cohorts included patient's age, gender, radiological fusion rate at 6months, radiological fusion rate at 12months and post-operative complications like cage subsidence, painful donor site, subluxation, limb pain, limb weakness, and limb numbness.

Results: Out of 52 patients, 90.4% achieved fusion at 12 months of follow up and 13.5% faced post-operative cage subsidence. The mean age of titanium cage group patients was 56.22 ± 15.93 years while that of PEEK cage group patients was 55.15 ± 16.11 years ($p=0.543$). The radiological fusion rate at 6 months was 62.5% in Titanium cage group and 60% in PEEK cage group ($p=0.543$). Similarly, fusion rate at 12 months follow up was 90.6% in first group and 90% in second group ($p=0.646$). Among post-operative complications, cage subsidence rate was 18.8% with Titanium cage and 5% with PEEK cage ($p=0.228$). One patient of PEEK cage group suffered painful donor site, one patient of Titanium group suffered subluxation and limb weakness and one patient of Titanium group suffered limb pain and numbness. Complications other than cage subsidence were also commutatively comparable in both cohorts ($p=0.673$).

Conclusion: The fusion rates with anterior cervical discectomy and fusion (ACDF) surgery are excellent and comparable using Titanium as well as PEEK cages with fewer post-operative complications. No one modality has superiority over other in our studied population. Titanium cages are cheaper, so may still be alternative to PEEK cages in our people where affordability is a big issue for general public.

Keywords: Anterior cervical discectomy and fusion (ACDF) surgery, Titanium cage, PEEK cage, SPSS

INTRODUCTION

Cervical spondylotic myelopathy (CSM)¹ is the consequence of cervical cord compression due to variety of spine disorders like spondylosis, trauma, neoplastic lesion, prolapsed disc and degenerative disc disease. Anterior cervical discectomy and fusion (ACDF)² is the procedure of choice for cervical spondylotic myelopathy and radiculopathy unresponsive to conservative care. This procedure was first reported in 1958 by Smith and Robinson.³ Its mortality rate is about 0.1%, and morbidity rate after its adverse events is 19.3%.^{4,5} The bony fusion rate after ACDF is excellent. In a meta-analysis of 146 articles having 10208 patients, fusion rate after one year of surgery was 90%.⁶

The complication rate associated with ACDF surgery varies from 0-4.3% for one level upto 56% for four level fusions,⁷ and include cage subsidence,⁸ dysphagia,⁹ vocal cord dysfunction,¹⁰ wound infection,¹¹ nerve root injury,¹² and CSF leak¹³ etc.

In ACDF procedure, discectomy is combined with interbody fusion. Cancellous bone graft is usually taken from the iliac crest. Many types of cages are used in ACDF

surgery for fusion in clinical practice. In our setup, titanium and polyetheretherketone (PEEK) cages¹⁴ are commonly used. The national as well as international data suggest no significant difference in outcome between Titanium and PEEK cages with a variable associated post-operative adverse events. In our setup, no such comparative study has been carried out. Therefore, the objective of this study was to compare the Titanium cage with PEEK cage in patients who underwent anterior cervical discectomy and fusion (ACDF) surgery at Neurosurgery Department of DHQ teaching hospital, Gujranwala.

MATERIAL AND METHODS


This was a retrospective analysis of the patients who underwent anterior cervical discectomy and fusion (ACDF) surgery from January 2015 to December 2018 at the Department of Neurosurgery, DHQ Teaching hospital, Gujranwala. The patients from both gender with more than two years follow up were included, while those with accompanying comorbidities like diabetes, end stage renal or lung disease, autoimmune disease, or underlying malignancy were excluded. The group of the patients among which titanium cage was used in ACDF surgery, were labelled as group A, while those in which PEEK cage was used, were labelled as group B. Informations

Received on 24-08-2019

Accepted on 18-01-2020

composed from both cohorts included patient's age, gender, radiological fusion rate at 6months, radiological fusion rate at 12months and post-operative complications like cage subsidence, painful donor site, subluxation, limb pain, limb weakness, and limb numbness. Postoperative imagings were obtained at 1, 6 and 12-months of follow-up for lucency around the work area and for issues with the cages like subsidence as well as post-operative complications. Cage subsidence⁸ was defined as equal or more than 3 mm loss of height comparing the direct postoperative intervertebral height with the intervertebral height at the last follow-up time. Adequate lucency i.e. upto 70-80% seen around screws on radiographs defined the union / fusion. All findings were recorded in a structured performa. SPSS Vr 25 was used. Means with standard deviations were computed of quantitative variables, and frequencies-percentages for qualitative variables. Chi-square test for independence and Independent sample T test were used for qualitative and quantitative variables respectively to determine their significant association with type of the cage used in ACDF surgery. The p values were taken statistically significant if < 0.05.


Among postop complications, cage subsidence rate was 18.8% with Titanium cage and 5% with PEEK cage (p=0.228). One patients of PEEK cage group suffered painful donor site, one patient of Titanium group suffered subluxation and limb weakness and one patient of Titanium group suffered limb pain and numbness. Complications other than cage subsidence were also commutatively comparable in both cohorts (p=0.673) (Table 2).



Picture 1: Fusion rate in patients who underwent ACDF surgery (n=52)

RESULTS

Out of 52 patients who underwent anterior cervical discectomy and fusion (ACDF) surgery, 90.4% achieved fusion at 12 months of follow up (Picture 1) and 13.5% faced host-operative cage subsidence (Picture 2). In 32 patients, Titanium cage while 20 patients PEEK cage was used in surgical protocol. Both cohorts had similar ages of the patients (mean age in group A 56.22 ± 15.93 vs mean age of group B 55.15 ± 16.11 years, p=0.543) (Table 1, 2).



Picture 2: Cage subsidence rate in patients who underwent ACDF surgery (n=52)

The radiological outcome in term of fusion and post-operative complications were comparable in both cage groups. The radiological fusion rate at 6 months was 62.5% in Titanium cage group and 60% in PEEK cage group (p=0.543). Similarly, fusion rate at 12 months follow up was 90.6% in first group and 90% in second group (p=0.646).

Table 1: Comparison of mean age with type of cage used in Anterior cervical discectomy & fusion (ACDF) surgery (n = 52)^{*}

Quantitative variables	Posterior pedicle screw fixation		Mean difference	p-value
	Titanium (mean+SD)	PEEK (mean+SD)		
Age of patients (years)	56.22±15.93	55.15±16.11	1.069	0.816

^{*}Independent sample T-test was used

Table 2: Comparison of various qualitative variables with type of cage used in Anterior cervical discectomy & fusion (ACDF) surgery (n = 52)

Predictors / Factors	Type of cage used in surgery		Total	p-value
	Titanium	PEEK		
Male	20 (62.5%)	12 (60%)	32 (61.5%)	0.543
Female	12 (37.5%)	8 (40%)	20 (38.5%)	
Fusion at 6 months:				0.543
Yes	20 (62.5%)	12 (60%)	32 (61.5%)	
No	12 (37.5%)	8 (40%)	20 (38.5%)	
Fusion at 12 months:				0.646
Yes	29 (90.6%)	18 (90%)	47 (90.4%)	
No	3 (9.4%)	2 (10%)	5 (9.6%)	
Cage subsidence:				0.228
Yes	6 (18.8%)	1 (5%)	7 (13.5%)	
No	26 (81.2%)	19 (95%)	45 (86.5%)	
Other complications:				0.673
Yes	2 (6.3%)	1 (5%)	3 (5.8%)	
No	30 (93.7%)	19 (95%)	49 (94.2%)	

Chi-square test for independence was used

DISCUSSION

Anterior cervical discectomy and fusion (ACDF) surgery has outstanding radiological fusion rates in cervical cord disease patients. Kevin Phan et al¹⁸ achieved a fusion rate of 96% after ACDF surgery where both titanium and PEEK cages were found equally effective. Yu-Cheng Chou¹⁶ observed 46.51% and 100% fusion rates in titanium and PEEK cage groups respectively. Mario Cabraja¹⁹ found a fusion rate 93.2% with titanium cage and 88% with PEEK cage. Similarly, in a meta-analysis by Zhi Jan-Li,¹⁷ yield was comparable with both kinds of cages.

In our study, the radiological fusion rate at 12 months follow up was 90.4%, which was 90.6% in titanium cage group and 90% in PEEK cage group. Hence, choice of the cage keeping in view the fusion rates can be either of titanium or PEEK cage, as both are equally effective.

Iris Noodhock and colleagues⁸ reviewed 71 studies including 4784 patients to determine cage subsidence rate among patients who underwent ACDF surgery. They found the mean incidence of subsidence of 21%. There was a little difference in subsidence rates depending on the material of the cage. In that study, subsidence rate with cage-screw-combination was 15.1%, with PEEK cage 23.5%, titanium cage 24.9%, and with polymethyl methacrylate (PMMA) cage was 30.2%.

In our study, after ACDF surgery, cage subsidence rate was 13.5% only. The subsidence in titanium cage group of the patients was 18.8% while in PEEK cage group was 5%, however correlation was insignificant ($p=0.228$). It means the choice of the cage is not affected by its subsidence rate in our patients.

There are numerous complications associated with ACDF other than subsidence. Tao Wang⁹ studied the dysphagia after ACDF surgery and observed a dysphagia rate of 5.4% and 0.4% at 3 and 24 months post-operative respectively while Kenneth Kam Leung Yeung¹⁰ studied 69 ACDF patients for post-procedural complications and found 2.9% hoarseness due to vocal cord dysfunction.

Wound infection is a common complication for all surgical procedures, however in a 8887 studied patients after ACDF,¹¹ post-operative wound infection rate was only 0.07%.

The iatrogenic nerve root injury may also occur in ACDF procedure which may manifest with limb pain, limb numbness, or limb weakness. Jason Pui Yin Cheung et al¹² explained the risk of nerve injury of about 2% during ACDF surgery.

The pain at the donor site is commonly seen after ACDF surgery. Silber and colleagues²⁰ reported 26.1% pain rate at donor site after this surgery and 11.2% chronically use analgesics for pain. M. Skeppholm et al²¹ observed that pain at donor site is a major issue only during first 4 weeks, after which it does not affect the quality of the life. In our study, only one patient suffered painful donor site, one faced subluxation and limb weakness and one suffered limb pain and numbness. In our study, cumulatively complications other than subsidence has a rate of only 5.8%, where titanium cage group has 6.3% and PEEK cage group has 5% rates with insignificant association between two groups ($p=0.673$).

Hence, considering fusion rate and complication rate including cage subsidence, no significant difference was found between patients of titanium and PEEK cage groups in our studied population.

CONCLUSION

The fusion rates with anterior cervical discectomy and fusion (ACDF) surgery are excellent and comparable using Titanium as well as PEEK cages with fewer post-operative complications. No one modality has superiority over other in our studied population. Titanium cages are cheaper, so may still be alternative to PEEK cages in our people where affordability is a big issue for general public.

REFERENCES

1. Wang B, Lu G. & Kuang L. Anterior cervical discectomy and fusion with stand-alone anchored cages versus posterior laminectomy and fusion for four-level cervical spondylotic myelopathy: a retrospective study with 2-year follow-up. *BMC Musculoskelet Disord* 2018; 19, 216.
2. Bano F, Zreik J, Alvi MA, Goyal A, Freedman BA, and Bydon M. Anterior Cervical Corpectomy and Fusion Versus Anterior Cervical Discectomy and Fusion for Treatment of Multilevel Cervical Spondylotic Myelopathy: Insights from a National Registry. *World Neurosurgery* 2019; 132: e852-e861.
3. Song KJ, Choi BY. Current concepts of anterior cervical discectomy and fusion: a review of literature. *Asian Spine J.* 2014;8(4):531-539.
4. Chen, Y., Zhang, L., Li, E. *et al.* Late deep cervical infection after anterior cervical discectomy and fusion: a case report and literature review. *BMC Musculoskelet Disord* 2019; 20: 437.
5. Fountas KN, Kapsalaki EZ, Nikolakakos LG, et al. Anterior cervical discectomy and fusion associated complications. *Spine (Phila Pa 1976)*. 2007; 32:2310-7.
6. Noordhoek, I., Koning, M.T. & Vleggeert-Lankamp, C.L.A. Evaluation of bony fusion after anterior cervical discectomy: a systematic literature review. *Eur Spine J* 2019; 28: 386-399.
7. Epstein NE. A Review of Complication Rates for Anterior Cervical Discectomy and Fusion (ACDF). *Surg Neurol Int.* 2019; 10:100.
8. Noordhoek I, Koning MT, Jacobs WCH, Vleggeert-Lankamp CLA. Incidence and clinical relevance of cage subsidence in anterior cervical discectomy and fusion: a systematic review. *Acta Neurochir (Wien)*. 2018;160(4):873-880.
9. Feng-Yu L, Da-Long Y, Wen-Zheng H, Li-Shuang H, Lei M, Hui W, et al. Risk factors for dysphagia after anterior cervical spine surgery: A meta-analysis. *Medicine* 2017; 96 (10): e6267.
10. Yeung KKL, Cheung PWH and Cheung JPY. Anterior cervical discectomy and fusion for cervical myelopathy using stand-alone tricortical iliac crest autograft: Predictive factors for neurological and fusion outcomes. *Journal of Orthopaedic Surgery* 2019; 27(3): 1-9.
11. Ghobrial GM, Harrop JS, Sasso RC, et al. Anterior Cervical Infection: Presentation and Incidence of an Uncommon Postoperative Complication. *Global Spine J.* 2017;7(1 Suppl):12S-16S.
12. Cheung JP, Luk KD. Complications of Anterior and Posterior Cervical Spine Surgery. *Asian Spine J.* 2016;10(2):385-400.
13. Weinheimer WJ, Rowley JG, Otto R, Floyd J, and Chen PG. Oropharyngeal CSF Leak Secondary to Anterior Cervical Discectomy and Fusion. *Case Reports in Otolaryngology* 2018; 1: 1-4.

14. Laratta JL, Reddy HP, Bratcher KR, McGraw KE, Carreon LY, Owens RK 2nd. Outcomes and revision rates following multilevel anterior cervical discectomy and fusion. *J Spine Surg.* 2018;4(3):496–500.
15. Junaid M, Rashid MU, Bukhari SS, Ahmed M. Radiological and clinical outcomes in patients undergoing anterior cervical discectomy and fusion: Comparing titanium and PEEK (polyetheretherketone) cages. *Pak J Med Sci.* 2018;34(6):1412–1417.
16. Chou Y, Chen D, Hsieh WA, Chen W, Yen P, Harnod Tet al. Efficacy of anterior cervical fusion: Comparison of titanium cages, polyetheretherketone (PEEK) cages and autogenous bone grafts. *Journal of Clinical Neuroscience* 2008; 15(11): 1240-1245
17. Li ZJ, Wang Y, Xu GJ, Tian P. Is PEEK cage better than titanium cage in anterior cervical discectomy and fusion surgery? A meta-analysis. *BMC Musculoskelet Disord.* 2016;17(1):379.
18. Phan, K., Pelletier, M.H., Rao, P.J., Choy, W.J., Walsh, W.R. and Mobbs, R.J. Integral Fixation Titanium/Polyetheretherketone Cages for Cervical Arthrodesis: Evolution of Cage Design and Early Radiological Outcomes and Fusion Rates. *Orthop Surg.*2019; 11: 52-59.
19. Cabraja M, Oezdemir S, Koeppen D, Kroppenstedt S. Anterior cervical discectomy and fusion: comparison of titanium and polyetheretherketone cages. *BMC Musculoskelet Disord.* 2012; 13:172.
20. Silber JS, Anderson DG, Daffner SD, Brislin BT, Leland JM, Hilibrand AS, et al. Donor site morbidity after anterior iliac crest bone harvest for single-level anterior cervical discectomy and fusion. *Spine* 2003;28(2):134-9.
21. Skeppholm M, Olerud C. Pain from donor site after anterior cervical fusion with bone graft: a prospective randomized study with 12 months of follow-up. *Eur Spine J.* 2013;22(1):142–147. doi:10.1007/s00586-012-2456-