

Frequency of Hyponatremia in Children with Pneumonia

MUHAMMAD AFZAL¹, MUAZZAM FUAAD², MUHAMMAD ANWAR³, AVINASH PUNSHI⁴, SANAULLAH⁵, SHAZIA SIDDIQUE⁶

¹Senior Registrar, Department of Medicine, Avicenna Medical College Lahore

²Assistant Professor, Department of Medicine, Doctor Trust Teaching Hospital/Rai Medical College, Sargodha

³Associate Professor, ⁵Assistant Professor, Department of Paediatrics, Rashid Latif Medical College Lahore

⁴Junior Consultant, MICU South City Hospital, Karachi

⁶Assistant Professor of Medicine, Fatima Jinnah Medical University Lahore

Correspondence to: Dr. Muhammad Afzal E-mail: dr.afzal.hassan@gmail.com 0313-0472424

ABSTRACT

Aim: To examine the frequency of hyponatremia in children with pneumonia.

Study Design: Cross-sectional

Place and Duration of Study: Department of Medicine, Avicenna Medical College Hospital, Lahore from 1st January 2020 to 30th June 2020.

Methodology: One hundred and eight children of both genders with ages 5 months to 15 years admitted for pneumonia were included. Patients detail demographics age, sex and severity of pneumonia were recorded. Blood sample was taken from the patients to examine serum sodium level. Hyponatremia was defined as serum sodium level less than 135mmol/L.

Results: Seventy one (65.74%) cases were males and 37 (34.26%) cases were females. Seventy three (67.59%) patients were ages ≤ 10 years while 35 (32.41%) were ages 11 to 15 years. 46 (42.59%) patients had severe pneumonia. Hyponatremia was found in 42 (38.89%) patients while 66 (61.11%) patients had serum sodium level >135 nmol/L. Hyponatremia was significant associated with severe pneumonia with p-value <0.05 .

Conclusion: Frequency of hyponatremia in children with pneumonia was high and was significantly associated with severe pneumonia.

Key words: Frequency, Hyponatremia, Pneumonia, Children

INTRODUCTION

Pneumonia in children under 5 years of age is the leading cause of death worldwide. Advances in the medicine sector, improved antibiotics and vaccines have decreased the incidence in developing countries of lower respiratory tract infection. However, pneumonia, which accounts for roughly 20% of infant deaths, is currently the leading cause of death in children under the age of 5 in developed countries¹, as pneumonia remains a major cause of morbidity and death in developing countries. In 2015, 178,717 children under the age of five died of pneumonia in India, the study, entitled "Fight for breathes," by Save the Children a non-profit organization, said that in 2015, twenty children died of the disease every hour in India that year.²⁻⁴

In the absence of timely therapy, the disease requiring hospitalization will progress. Pneumonia-admission kids are seriously ill, and complications frequently arise, including electrolyte irregularities, most commonly hyponatremia. Studies in western countries found that hyponatremia was present in up to 45.4% of children hospitalized by pneumonia.^{3,5} Pneumonia hyponatremia was associated with an excessive anti-diuretic hormone (SIADH) secretion syndrome.⁶ Excess ADH results in the accumulation of water and expansion of volume, which contribute to serum osmolality decreasing below the reference level. Hyponatremia occurs only when the patient eats or gets some free water supply. In most pneumonia children, fluid therapy is not needed as a result of breathlessness, tiredness, or the risk of aspiration.⁷

The retention of water due to the secondary SIADH degradation of renal water has been identified in several pulmonary conditions, including in children infected by less respiratory tract infection.⁸ The current research was thus conducted to detect the occurrence and potential impact of

hyponatremia on children between 5 months and 15 years of age as well as the consequence of extreme pneumonia.

MATERIALS AND METHODS

This cross sectional study was conducted at Department of Medicine, Avicenna Medical College Hospital Lahore from 1st January 2020 to 30th June 2020. A total of 108 patients of both genders presented with pneumonia were enrolled. Patient's ages were ranging between 5 months to 15 years were included. Detailed demographics including age, sex, severity of disease and complete blood count were recorded. Children with other serious conditions were removed from the study such as heart disease, renal disease, and those with no consent were excluded. 2 ml blood sample was taken from each patient and sent to laboratory to examine serum sodium level. Serum sodium level <135 nmol/L was defined as hyponatremia. Hyponatremia was considered as mild when sodium level between 130-134 nmol/L, between 125-129 nmol/L considered as moderate and <125 nmol/L considered as severe. Association of hyponatremia with severity of disease was examined. All the data was analyzed by SPSS 24. Chi-square test was done to stratified hyponatremia by disease severity. P-value <0.05 was taken as significant.

RESULTS

Out of 108 patients, 71 (65.74%) were males and 37 (34.26%) were females in this study. 73 (67.59%) patients were ages ≤ 10 years while 35 (32.41%) were ages 11 to 15 years. 46 (42.59%) patients had severe pneumonia (Table 1). Hyponatremia was found in 42 (38.89%) patients while 66 (61.11%) patients had serum sodium level >135 nmol/L (Fig. 1). When we stratified hyponatremia with pneumonia

and severe pneumonia, we found that frequency of hyponatremia was significantly high in patients with severe pneumonia 32/46 (59.57%) as compared to patients with pneumonia 10/62 (16.13%), the difference was statistically significant [P<0.0001] (Table 2).

Table 1: Baseline details of patients

Variable	No.	%
Age (years)		
<10	73	67.59
11 – 15	35	32.41
Gender		
Male	71	65.74
Female	37	34.26
Severe pneumonia		
Yes	46	42.59
No	62	57.41




Fig. 1: Frequency of hyponatremia among all the patients

Table 2: Association of hyponatremia with disease severity

Variable	Disease Severity		Total
	Pneumonia	Severe Pneumonia	
Hyponatremia	10 (16.13%)	32 (59.57%)	42
No Hyponatremia	52 (83.87%)	14 (40.43%)	66
Total	62	46	108

P<0.0001

DISCUSSION

One of the most popular electrolyte abnormalities in hospitalized patients has been hyponatremia. It complicates many conditions involving respiratory, cardiovascular and central nervous systems etc. and is normal in pneumonia children. Not only that, it can function as a sign of serious disease that contributes to high death and morbidity.^{9,10} In the present study, majority 65.74% patients were males while females 34.26%. Most of the patients 67.59% were ages between 5 months to 10 years while 32.41% patients were ages 11 to 15 years. These results were comparable to many of previous studies in which male children were high in numbers 55%-65% and majority of patients were ages less than 10 years.^{11,12}

In present study we found that 46 (42.59%) patients had severe pneumonia while 57.41% patients had no severe pneumonia. A study conducted by Praneetha et al¹³ regarding hyponatremia in hospitalized children with pneumonia and they reported that out of 122 children 54 (44.26%) patients had severe pneumonia according to the WHO classification of pneumonia.

In the present study, hyponatremia was found in 42 (38.89%) patients while 66 (61.11%) patients had serum sodium level >135 nmol/L. Mandal et al¹⁴ demonstrated that hyponatremia was found in 21% out of 100 pediatric patients of pneumonia. Another study conducted by Jha et al¹⁵ reported that hyponatremia was highly associated with pneumonia and the frequency of hyponatremia among children with pneumonia was 80%.

Praneetha et al¹³ reported that hyponatremia was found in 43.44% patients presented with pneumonia. Some other studies demonstrated that the frequency of hyponatremia accounted 32% to 55% among children with pneumonia.^{16,17}

In our study, the frequency of hyponatremia was significantly high in patients with severe pneumonia 32/46 (59.57%) as compared to patients with pneumonia 10/62 (16.13%), the difference was statistically significant with p-value <0.0001. These results showed similarity to many of previous studies in which children with severe pneumonia were more prone to have hyponatremia as compared to children with pneumonia.^{14,18}

Complication incidence is about three-fold in children with hyponatremia as compared to normonatremic patients. Studies also have shown that a comorbidity factor in respiratory compromises is substantially increased pneumonia deaths in patients with hyponatremia.^{19,20} The majority of patients reported mild pneumonia in 63.44% and moderate pneumonia in 21.5% and 15.5% children had severe hyponatremia.

CONCLUSION

Electrolyte abnormalities are the leading cause of increasing morbidity and mortality in children with severe disease. Hyponatremia, is one of the main disorders that affect favourable outcomes in children with pneumonia. We concluded that frequency of hyponatremia in children with pneumonia was high and was significantly associated with severe pneumonia.

REFERENCES

- Zar HJ, Madhi SA. Childhood pneumonia-progress and challenges. South Afr Med J 2006; 96(9):890-9.
- Save the children. Fighting for breath. Available at: <https://www.savethechildren.org.uk/content/dam/global/reports/health-and-nutrition/fighting-for-breathlow-res.pdf>. Accessed 12 January 2018
- Subba RSD, Thomas B. Electrolyte abnormalities in children admitted to pediatric intensive care unit. Indian Pediatr 2000; 37(12):1348-53.
- Black RE, Cousens S, Johnson HL. Global, regional, and national causes of child mortality in 2008: a systematic analysis. Lancet 2012; 375(9730):1969-19.
- Swart RM, Hoorn E J, Betjes MG, Zieste R. Hyponatremia and inflammation: the merging role of interleukin-6 in osmoregulation. Nephron Physiol 2011; 118(2):45-51.
- Asadollahi K, Beeching N. Gill Hyponatremia as a risk factor for hospital mortality. Q J Med 2006;99:877-80
- Hawkins RC. Age and gender as risk factors for hyponatremia and hypernatremia. Clin Chim Acta 2003;337:169-72
- Nair V, Niederman MS, Masani F, Fishbane S. Hypo natremia in community-acquired pneumonia. Am J Nephrol 2007;27:184-90.

9. Zilberberg MD, Exuzides A, Spalding J, Foreman A, Jones AG, Colby C, et al. Hyponatremia and hospital outcomes among patients with pneumonia: a retrospective cohort study. *BMC Pulmonol Med* 2008;8(1):16
10. Wrotek A, Jackowska T. Hyponatremia in children hospitalized due to pneumonia. *Adv Exp Med Biol* 2013;788:103-8
11. Müller M, Schefold JC, Guignard V, Exadaktylos AK, Pfortmueller CA. Hyponatraemia is independently associated with in-hospital mortality in patients with pneumonia. *Eur J Int Med* 2018;54:46-52
12. Tagarro A, Martín MD, Del-Amo N, Sanz-Rosa D, Rodríguez M, Galán JC, et al. Hyponatremia in children with pneumonia rarely means SIADH. *Paediatr Child Health* 2018;23(7):e126-33.
13. Praneetha CK, Ahirrao VSK, Srinivasa R, Ravichander RB. Hyponatremia in children of 2 months to 5 years of age with community acquired pneumonia and its correlation with severity of illness and outcome. *Int J Pediatr Res* 2019; 6(11): 561-6.
14. Mandal PP, Garg M, Choudhary IP. To study the association and significance of hyponatremia in pneumonia in paediatric patients treated in hospital setting. *Int J Contemp Med Res* 2018;5(1):11-4.
15. Jha C, Tamrakar A. Assessment of hyponatremia in pneumonia in children. *Br J Health Sci* 2019; 3(3): 542-7.
16. Park SW, Shin SM, Jeong M, Cho DH, Lee KH, Eisenhut M, et al. Hyponatremia in children with respiratory infections: a cross-sectional analysis of a cohort of 3938 patients. *Sci Rep* 2018;8: 16494.
17. Don M, Valerio G, Canciani M, Korppi M. Hyponatremia in radiologically confirmed pediatric community-acquired pneumonia. *Pediatr Emerg Care* 2014;30(1):86.
18. Das M, Narain B. Hyponatremia in children with severe pneumonia and its effect on overall outcome. *Int J Contemp Pediatr* 2019;6:2516- 21.
19. Kumar K, Devi A, Sharma RK. Hyponatremia in children of 2 months to 5 years of age with pneumonia and its correlation with outcome. *Int J Med Biomed Studies* 2020; 4(5).
20. Sakellaropoulou A, Hatzistilianou M. Hyponatremia in cases of children with pneumonia. *Arch Med Sci* 2010; 4: 578–83.