

Antidiabetic effect of Aqueous Extract of *Medicago Sativa* with Enhanced Histopathology of Pancreas in Alloxan Induced Diabetic Rats

SALIM J.KHALAF¹, GADEER HATEM ALJADER¹, ENTEDHAR R. SARHAT^{1*}, THURAIYA RIFAAT SARHAT², AND KASIM SAKRAN ABASS³

¹Department of Basic science, Dentistry College, University of Tikrit, Tikrit, Iraq

²College of Education, University of Tikrit, Tikrit, Iraq

³Department of Pharmacology and Toxicology, College of Pharmacy, University of Kirkuk, Kirkuk, Iraq

Corresponding author to Dr. Entedhar R. Sarhat entedharr@tu.edu.iq

ABSTRACT

Aim: The present study investigated the effect of *Medicago sativa* extract and glibenclamide on biochemical parameters in the liver as well as on pancreas tissue in alloxan-induced diabetic rats.

Methods: Diabetes mellitus was induced in 28 out of 35 adult male albino rats, using an intra-peritoneal injection of 65 mg/kg body weight of alloxan. The diabetic rats were divided into four groups, two of which were administered orally by garlic extract (250 and 500 mg/kg) and a group composed of diabetic rats was given the standard drug, glibenclamide, orally at a dose of 2.5 mg/kg. The control rats (normal and diabetic) were fed normal saline, once daily for 21 d. The specimens were prepared for light microscopic examination. In parallel, the related biomedical parameters such as glucose and insulin levels had been estimated in pancreas, statistically analyzed and compared between the groups.

Results: The aqueous extract of *M. Sativa* significantly reversed ($P < 0.05$) the manifestation of alloxan on the levels of serum glucose & insulin, C-peptide, malondialdehyde formation, Nitric oxide, glucose-6-phosphate dehydrogenase (G6PD), antioxidant enzymes (superoxide dismutase, glutathione peroxidase and catalase). Histological pictures of pancreas showed pathological changes which was in concurrence with the biochemical results.

Conclusion: Extracts of *Alfalfa* improved hyperglycemia and other biochemical alterations noticed in alloxan-diabetic rats. These effects may be due to the presence of a high content of flavonoids which acts synergistically as antioxidants.

Keywords: Diabetes mellitus ; Pancreas; *M. Sativa*; Glibenclamide.

INTRODUCTION

Pancreas is a vital endocrine-exocrine organ that produces several hormones and enzymes. Its enzymes help in the digestion of carbohydrates, fats, and proteins whereas its hormones such as insulin regulate carbohydrate metabolism in the body and maintains passage of glucose across the cell membrane. Therefore, any change in the function of the organ may directly affect the physiological function of the body¹. Diabetes mellitus (DM), an endocrine metabolic disorder of multiple etiologies manifested by consistent elevated levels of glucose in the blood resulting from defects in insulin secretion, insulin action, or both, and occurs in almost all populations of the world a variable prevalence^{2,3}. Recently there has been a growing interest in alternative therapies, including the use of plant foods, to treat diabetic patients⁴.

Alfalfa (Medicago sativa) or green gold is a high-yielding perennial legume that is cultivated worldwide with rich nutritional characteristics and bioactive compounds, that are used in traditional medicine due to being high in protein, calcium, β -carotene vitamins, including (B, C, E, and K), chlorophyll, c omarine derivative, choline, essential amino acid, flavonols, lime, magnesium, phosphorous, protein, silicon, potassium, sterol, iron and Saponins and low in percentage of cellulose, lignin, and xylans,. It contains many enzymes, including amylase, invertase, and pectinase, and so it can be used as digestive aids⁵⁻⁷.

Glyburide (glibenclamide) is one of the sulfonylurea compound chemically, 5-chloro-N- [2- [4-cyclohexyl carbamoyl sulfamoyl) phenyl] ethyl]-2-methoxy benzamide, to be good alternative for insulin. Oral hypoglycemic agents are an attractive option to insulin because of their lower cost and ease of administration, which increase patient compliance which is produced its effect via stimulating of endogenous insulin release from pancreas, enhancing peripheral tissue utilization of glucose, or by decreasing the absorption of glucose by intestine⁸⁻¹⁰.

This study aimed at testing the ameliorative effect of two low doses of *Moringa* seeds powder (50 and 100 mg/kg body weight) on alloxan induced diabetic male rats.

MATERIALS AND METHODS

Induction of Diabetes: Alloxan is known as a useful drug to induce diabetes in experimental animals. After overnight fasting diabetes was induced in rats by intraperitoneal injection of alloxan monohydrate (Sigma, St. Louis, MO, USA) dissolved in 1ml distilled water at a dose of 65mg/kg BW.

Preparation of *Medicago sativa* Extract: Leaves of *Medicago sativa* after collection were allowed to shade dry and the dried leaves were ground to a fine powder, which was then used for extraction in a soxhlet apparatus for up to 10 hours using 95% ethanol at a boiling temperature of 60°C. The extract obtained from soxhlet extraction was

allowed to cool and then filtered to remove the residue. The filtrate was then concentrated at 65°C by rotavapour to get a fine powder that was refrigerated at 4°C until further use¹¹.

Study design: A total of 35 Male albino Wistar rats of age 4-5 weeks with a body weight of 180 to 200g were procured and acclimatized for 7 days to animal house and were randomly divided into 5 groups each consisting of 5 rats as follows:

Group 1: Non-diabetic control-received distilled water (3ml/kg.b.w).

Group 2: Diabetic control- received 1 ml of distilled water

Group 3: Diabetic - received Medicago sativa extract (60mg/kg.b.w).

Group 4: Diabetic - received Medicago sativa extract (120mg/kg.b.w).

Group 5 : Diabetic - received Glibenclamide (10 mg/kg.b.w).

They were housed and received normal basal diet and tap water *ad libitum* in a constant environment (room temperature 28 ± 2°C, room humidity 60 ± 5%) with a 12/12 h light/ dark cycle.

Medicago sativa and glibenclamide doses were administered daily via oral gavage. Food and water intake were monitored daily whilst body weights were determined weekly.

Insulin levels were measured by Enzyme-linked Immunosorbent Assay (ELISA) technique. G6PDH activity was assayed by the method of Beutler. Catalase, MDA, GPx, G6PD, NO levels were measured by spectrophotometric kit.

At the end of the experimental period the rats were sacrificed using ether anesthesia. pancreas were dissected, fixed in 10% buffered formalin, and then dehydrated by successively passing through a gradient mixture of ethanol and water. The samples were rinsed in xylene and embedded in paraffin. 5 µm thick sections were prepared and stained with hematoxylin and eosin (H&E) dye for microscopic investigation. The stained sections were examined and photographed under a light microscope^{12,13}.

Statistical analysis :One-way-ANOVA-using- SPSS-statistics. The p-value of more than 0.05 was considered as not significant.

RESULTS

Pancreatic insulin SOD, GPx, G6PD, NO and catalase levels were significant ($P < 0.05$) decreased while MDA level was significant increase in diabetic control rats as compared with normal control rats.

Treatment with aqueous extract of *M. Sativa* significantly enhanced the activity of SOD, GPx, G6PD, NO and catalase while MDA level decreased. Glibenclamide treatment produced significant increases in SOD, GPx, G6PD, NO and catalase while MDA decreased, compared to the diabetic control group.

Figure.1: Effect of *M. Sativa* on serum insulin levels in diabetic rabbits.




Figure.2: Effect of *M. Sativa* on serum GPx levels in diabetic rabbits




Figure.3: Effect of *M. Sativa* on serum catalase levels in diabetic rabbits




Figure.4: Effect of *M. Sativa* on serum NO levels in diabetic rabbits




Figure.5: Effect of *M. Sativa* on serum G6PD levels in diabetic rabbits




Figure.6: Effect of *M. Sativa* on serum MDA levels in diabetic rabbits




Table 3 - Effect of the Orally Administered *M. Sativa* Extract and Glibenclamide on Biochemical Parameters .

Parameters	Control	Group B	Group C	Group AD	Group E
Insulin (ng/ml)	2.2 ± 0.078	0.9 ± 0.059***	1.9 ± 0.08	1.51 ± 0.032**	2 ± 0.087
GPx (mU/ml)	7 ± 0.24	1.9 ± 0.56	4.43 ± 0.75#	6 ± 0.46	5.82 ± 0.85
Catalase (mU/ml)	0.42 ± 0.09	0.2 ± 0.03*	0.47 ± 0.06#	0.45 ± 0.09#	0.59 ± 0.08#
NO (uM)	35.42 ± 1.17	22 ± 0.95**	29.2 ± 1.65#	33 ± 3.4#	31 ± 3.7#
G6PDIU/g HB	5.62 ± 0.05	2.75 ± 0.13	4.82 ± 0.11	5.24 ± 0.17	4.46 ± 0.09
MDA	2.29 ± 0.23	4.03 ± 0.23	2.40 ± 0.45**	2.15 ± 0.26**	2.13 ± 0.20 a

*Significantly different from non-diabetic control p < 0.05

**Significantly different from non-diabetic control p < 0.01

*** Significantly different from non-diabetic control p < 0.001

#Significantly different from diabetic control p < 0.05

##Significantly different from diabetic control p < 0.01

DISCUSSION

The current results reveals significant decrease in levels of serum insulin and marked elevation in levels of blood glucose in diabetic rats. This is attributed to the hypo secretion of insulin by the pancreatic β-cells, as alloxan selectively destroys the pancreatic insulin secreting β-cells and induces hyperglycemia (14).The current observations are in analogy to earlier results obtained by (15,16).The concurrent treatment with alfalfa these parameters and nearly restored them to their normal levels. This curative effect is due to the active constituents present in alfalfa as trigonelline, which is the N-methyl derivative and main human metabolite of the vitamin nicotinic acid¹⁷.

Hyperglycemia is the clinical hallmark of poorly controlled diabetes, which is known to cause protein glycation, also known as nonenzymatic glycosylation. It has been reported that various proteins, including

hemoglobin, albumin, collagen, low-density lipoprotein, a crystalline fibronectin, undergo non-enzymatic glycation in diabetes¹⁸.

Hence, estimation of glycosylated hemoglobin is a well-accepted biochemical parameter useful for the diagnosis and management of the disease. The increased glycated hemoglobin is associated with loss of β-cell function and has been implicated in the complications of diabetes mellitus^{19,20}.

The present study indicated significant increases in the activity of HbA1c in the STZ-induced diabetic rats and these levels were significantly reduced after treatment intervention with *Medicago sativa* which are concomitant with other findings²¹⁻²³.

Oxidative stress consider as interrelated contributing factors in pancreatic β-cell dysfunction and apoptosis^{24,25}. Because of poor antioxidant capacity, beta cells are

vulnerable to the oxidative stress induced by DM glucotoxicity (26).

Elevated level of MDA which considered one of the final products of lipid peroxidation (27). as well as reduced levels of key antioxidant enzymes, CAT and GPx levels in the pancreas of the diabetic control rats (Table 3), an indication of alloxan-induced oxidative stress. Glutathione is the mother of all antioxidants, the master detoxifier and maestro of the immune system. The low activity of GPx could be directly explained by the low content of GSH, since GSH is a substrate and cofactor of GPx. Enzyme inactivation could also contribute to low GPx activity²⁸⁻³⁰. A decrease in the activity of CAT could be due to increase in the lipid peroxidation product, malondialdehyde which can form cross links, thereby inactivating several membrane bound enzymes^{31,32}.

Treatment with alfalfa (60 and 120mg/kg) produced significant reductions in MDA as well as increased CAT and GPx levels demonstrating that the alfalfa leaves contain high concentrations of flavonoids and other phenols, which exert powerful antioxidant properties and capable of stimulating liver antioxidant enzymes as free radical scavengers and inhibitors of nitric oxide release and therefore may be capable of preventing tissue damage, it can protect the pancreas tissues from lipid peroxidation on diabetic rats (33,34).

Glucose-6-phosphate dehydrogenase (E.C.1.1.1.49) is the rate-limiting enzyme in the pentose-phosphate pathway, found in the mitochondria and cytosol. it converts glucose-6-phosphate accompanied by nicotinamide adenine dinucleotide phosphate NADP⁺ into 6-phosphogluconolactone and NADPH (Omoso,2018). G6PD 6PD provides a source of reducing power against oxidative damage and is important for β -cell proliferation and prevention of β -cell death³³⁻³⁵.

In alloxan-induced diabetic rats, the reduction of G-6-PDH activity in liver which obstruct glucose utilization through pentose phosphate pathway as this enzyme activity is controlled by insulin^{33,36}, and due to insufficient or limited production of NADPH which regenerates GSH, a physiologic antioxidant to scavenge glucose-generated free radicals²³.

The G6PDH activity was restored to normal level after treatment with *Medicago sativa* for 30 d. The restored G6PDH activity by *Medicago sativa* treatment confirmed the protective role of *Medicago sativa* against diabetes complications, since *Medicago sativa* administration showed a similar upturn with standard anti-diabetic drug glibenclamide.

In conclusion, *Medicago sativa* have potent anti-hyperglycemic and efficacies in alloxan-induced diabetic rats. These effects may be due their ability to improve the islets architecture, insulin secretory response and antioxidant activity.

Conflict of Interests: The authors of this paper declare that he has no financial or personal relationships with individuals or organizations that would unacceptably bias the content of this paper and therefore declare that there is no conflict of interests.

Source of Funding: The authors have no sources of funding, so it is self-funding research.

Ethical Approve: We declare that the study does not need ethical approval.

REFERENCES

1. Maisa MA Al-Qudah, Moawiya A Haddad, Jafar MF EL-Qudah. The effects of aqueous ginger extract on pancreas histology and on blood glucose in normal and alloxan monohydrate-induced diabetic rats. *Biomedical Research* 2016; 27 (2): 350-356
2. Sarhat ER, Rmaid ZJ, Jabir TH. Changes of salivary interleukine17, Apelin, Omentin and Vaspin levels in normal subjects and diabetic patients with chronic periodontitis, *Ann Trop Med & Pub Health*; 23:S404. DOI: <http://doi.org/10.36295/ASRO.2020.23118>
3. Rajesh R., S. Arunchandra Singh, K. Anandraj Vaithy, K. Manimekalai, Dhananjay Kotasthane, S. S. Rajasekar. THE EFFECT OF MUCUNA PRURIENS SEED EXTRACT ON PANCREAS AND LIVER OF DIABETIC WISTAR RATS. *IJCRR*.2016;(08) 04: 61-67
4. Entedhar R. Sarhat, Husamuldeen Salim Mohammed Saeed .Effects of Lycopene on Paraoxonase and Adipokines Parameters in Streptozotocin - Induced Diabetic Rabbits. *SUST Journal of Natural and Medical Sciences (JNMS)*. 18 (1); 2017:1-8.
5. Sayran Sattar Saleh , Entedhar Rifaat Sarhat. Effects of Ethanolic Moringa Oleifera Extract on Melatonin, Liver and Kidney Function Tests in Alloxan-Induced Diabetic Rats . *Indian Journal of Forensic Medicine & Toxicology*, 2019; 13(4): 1015- 1019.
6. Farsani MK, Amraie E, Kaviani P, Keshvari M. Effects of aqueous extract of alfalfa on hyperglycemia and dyslipidemia in alloxan-induced diabetic Wistar rats. *Interv Med Appl Sci*. 2016;8(3):103–108.
7. C. Fu, T. Hernandez, C. Zhou and Z. Y. Wang, *Alfalfa (Medicago sativa L.)*, *Agrobacterium Protocols*, Springer, New York, 2015: 213–221.
8. L. X. Yu , X. Liu , W. Boge and X. P. Liu , Genome-wide association study identifies loci for salt tolerance during germination in autotetraploid alfalfa (*Medicago sativa L.*) using genotyping-by-sequencing, *Front. Plant Sci.*, 2016;7 :956.
9. Mahmoud Bahmani, Mahmoud Mirhoseini, Hedayatollah Shirzad, Mehrmoosh Sedighi, , Nejme Shahinfard, and Mahmoud Rafieian-Kopaei. A Review on Promising Natural Agents Effective on Hyperlipidemia. *Journal of Evidence-Based Complementary & Alternative Medicine*.2015; 20(3) 228-238.
10. Shital S. Chavan, Ravindra S. Jadhav, Kavita S. Khemnar and Vishal B. Tambe. Evaluation of Antibacterial Activity and Phytochemical Screening of *Medicago sativa* Leaves . *Int.J.Curr.Res.Aca.ev*.2015; 3 (5):: 308-313.
11. Entedhar R. Sarhat , Ayhan R. Mahmood , Mohammed Sh. Abdulla. Effect of Glibenclamide and Tomato lycopene extract on some biochemical parameters in serum of alloxan Induced diabetic rabbits. *Kirkuk University Journal /Scientific Studies (KUJSS)*.2015; 10(3):140-154.
12. Zohar Nachum, Noah Zafran, Raed Salim, Noura Hissin, Jamal Hasan ein, Yifat Gam Ze Letova, Abeer Suleiman, Enav Yefet. Glyburide Versus Metformin and Their Combination for the Treatment of Gestational Diabetes Mellitus: A Randomized Controlled Study. *Diabetes Care* Mar 2017, 40 (3) 332-337.
13. Seena T. Pandarekandy, P. G. Sreejesh, B. S. Harikumar Thampi and E. Sreekumar. Hypoglycaemic Effect of Glibenclamide` ZQ: A Critical Study on the Basis of Creatinine and Lipid Peroxidation Status of Streptozotocin-induced Diabetic Rat. *Indian J Pharm Sci*. 2017;79(5):768-777.
14. Baxi, D. B., Singh, P. K., Doshi, A. A., Arya, S. Mukherjee, R. and Ramachandran A.V. *Medicago Sativa* leaf extract supplementation corrects diabetes induced dyslipidemia, oxidative stress and hepatic renal functions and exerts antihyperglycaemic action as effective as Metformin. *Annals of Biological Research*, 2010. 1 (3):107-119
15. Al-Malki AL, El Rabey HA. The antidiabetic effect of low doses of *Moringa oleifera* Lam. seeds on streptozotocin induced diabetes and diabetic nephropathy in male rats. *Biomed Res Int*. 2015;2015:381040. 1-13.
16. Buthayna A. A., Entedhar R. S. , Siham A. W. Study of The Effect of Castor Seeds(*Ricinus Communis* linn.)on Ovary Functions and Characters of Female Rabbits. . *Assiut Vet. Med. J*. 2017; 63 (152):62-65.
17. Eman G. E. Helal, Nouran Abou-Aouf1, Sayda M. Khattab, Hassan S. Shaibah, Hoda M. Abu-Taleb, Tamer M. M. Abu-Amara. The Effects of *Jasania montana* (Neheda) on Some Biochemical and Histological Parameters of Diabetic Albino Rats. *The Egyptian Journal of Hospital Medicine*.2014. 57:513-530.
18. Entedhar R. S. Effect of Ginger on the activity of some antioxidant enzymes (Superoxide dismutase, and Catalase) of Alloxan

- Experimental Induced-Diabetic Rabbits. *Al-Mustansiriyah J. Sci.* 2011; 22(5):192-200.
19. Entedhar R. Sarhat, Buthyna A. Abdullah, Siham A. Wadi. Experimental Study of Effect of Ginger Ethanolic extract on Oxidative Stress in Alloxan Induced-Diabetic Rabbits. 3rd Scientific Conference - College of Veterinary Medicine - University of Tikrit, 2016:35-38.
 20. Eman G. E. Helal, Samia M. Abd-Elwahab, Tarek A. Atia and Anwaar Alkamel Mohammad. Hypoglycemic Effect of the Aqueous Extracts of *Lupinus albus*, *Medicago sativa* (Seeds) and Their Mixture on Diabetic Rats. *The Egyptian Journal of Hospital Medicine.* 2013; 52:685-698.
 21. Iliya I A, Mohammed B, Akuyam S A, Yaro J D, Timbuak J A, Tanko M, Nok A J. Histological and biochemical evaluation of the antidiabetic potentials of s-allyl-cysteine and mangiferin in type 2 diabetic rat models. *Sub-Saharan Afr J Med.* 2016;3:32-40.
 22. LILIA DOUAOUYA, NOUREDDINE BOUZERNA. EFFECT OF GARLIC (*ALLIUM SATIVUM* L) ON BIOCHEMICAL PARAMETERS AND HISTOPATHOLOGY OF PANCREAS OF ALLOXAN-INDUCED DIABETIC RATS. *Int J Pharm Pharm Sci*, 2016; 8(6):202-206.
 23. Hui YangXun Jin Christopher Wai Kei Lam Sheng-Kai Yan. *Clinical Chemistry and Laboratory Medicine.* 2011;49(11):1773-82.
 24. Entedhar Rifaat Sarhat, Siham A. Wadi, Ban I Sedeeq. Thuraia R. Sarhat, Nawar.A. Jasim. Study of histopathological and biochemical effect of *Punica granatum* L. extract on streptozotocin -induced diabetes in rabbits. *Iraqi Journal of Veterinary Sciences*, 2019; 33(1):189-194.
 25. HOSSEINI, Azar; SHAFIEE-NICK, Reza and GHORBANI, Ahmad. Pancreatic beta cell protection/regeneration with phytotherapy. *Braz. J. Pharm. Sci.* 2015; 51(1): 1-16.
 26. Intesar Jasim Mohammed , Entedhar Rifaat Sarhat , Marwa Abdul-Salam Hamied ,Thuraia Rifaat Sarhat. Assessment of salivary Interleukin (IL)-6, IL-10, Oxidative Stress, Antioxidant Status, pH, and Flow Rate in Dental Caries Experience patients in Tikrit Province. *Sys Rev Pharm* 2021;12(1):55-59.
 27. Gawlik K, Naskalski J. W. Fedak, D., Pawlica-Gosiewska D., Grudzień U., Dumnicka P., Malecki M. T., Solnica B., "Markers of Antioxidant Defense in Patients with Type 2 Diabetes", *Oxidative Medicine and Cellular Longevity*, vol. 2016, ArticleID 2352361, 6 pages, 2016. <https://doi.org/10.1155/2016/2352361>
 28. Khalid G. Washeel, Entedhar R. Sarhat , Talal H. Jabir. Assessment of Melatonin and Oxidant-Antioxidant Markers in Infertile Men in Thi-Qar Province. *Indian Journal of Forensic Medicine & Toxicology*, 2019; 13(4):1500-1504.
 29. Entedhar R. Sarhat, Siham A. Wadi, Mutaz S. Ahmed, Shaima N. Mustafa, Thuraia R. Sarhat. Evaluation of Serum Malondialdehyde, Glutathione peroxidase, Superoxide dismutase, and Catalase levels in Hormonal Contraceptives. *Tikrit Medical Journal.* 2018; 24(1) :10-20.
 30. Karimi E, Oskoueian E, Oskoueian A, Omidvar V, Hendra R, Nazeran H. Insight into the functional and medicinal properties of *Medicago sativa* (alfalfa) leaves extract. *Journal of Medicinal Plants Research.* 2013;7:290-297.
 31. Omotoso G.O. , Adunmo G.O., Ojulari L.S., Olawuyi T.S., Lewu F.S., Jaji-Sulaimon R. , Sulaimon F.A., Gbadamosi I.T., Onoja O.P. Moringa oleifera attenuates biochemical and histological changes associated with the pancreas in nicotine-treated rats. *Res. J. of Health Sci.* 2018; 6(4):172-181.
 32. Zhang Z, Liew CW, Handy DE, et al. High glucose inhibits glucose-6-phosphate dehydrogenase, leading to increased oxidative stress and beta-cell apoptosis. *FASEB J.* 2010;24(5):1497-1505.
 33. Ahmed D, Kumar V, Verma A, Shukla GS, Sharma M. Antidiabetic, antioxidant, antihyperlipidemic effect of extract of *Euryale ferox* salisb. with enhanced histopathology of pancreas, liver and kidney in streptozotocin induced diabetic rats. *Springerplus.* 2015;4:315.
 34. Engwa GA, Nwalo FN, Chibuzor GE, Ejiagha EC, Abonyi MC, et al. Relationship between Type 2 Diabetes and Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency and Their Effect on Oxidative Stress. *J Diabetes Metab.* 2018; 9: 800.
 35. Mahde S. Hamad, Entedhar R. Sarhat, Salim J. Khalaf, Thuraia Rifaat Sarhat, Kasim Sakran Abass . 2021. Characteristic Abnormalities In Serum Biochemistry In Patients With Breast Cancer. *Systematic Reviews in Pharmacy*, 11(11): 1967-1971.
 36. Zubaida N.M Albarzanji, Thikra Abdullah Mahmood, Entedhar Rifaat Sarhat , Kasim Sakran Abass. 2020. Cytokines Storm Of COVID-19 And Multi Systemic Organ Failure: A Review. *Systematic Reviews in Pharmacy* 11(10), 1252-1256