

# Deficiency of Vitamin D Levels in First Trimester of Pregnancy Among Primigravida Women

AROJ FATIMA<sup>1</sup>, SHEHNAZ SHEEBA<sup>2</sup>, NAVEEN IQBAL<sup>3</sup>, AESHA SADAF RIZWAN<sup>4</sup>, UMBER ASAD<sup>5</sup>

<sup>1</sup>FCPS Gynae & Obs, Senior Registrar Amna Inayat Medical College, Lahore

<sup>2</sup>MCPS MRCOG, Gynaecologist, PAF Hospital E-9 Islamabad

<sup>3</sup>Senior Registrar Gynae & Obs, CMH Kharian Medical College, Kharian

<sup>4</sup>Assistant Professor of Obstetrics and Gynaecology, Shahida Islam Medical College, Lodhran

<sup>5</sup>Senior Registrar Gynae & Obs, Unit-II Sir Ganga Ram Hospital Lahore

Corresponding Author: Dr Arooj Fatima, Email: amnafiza22@gmail.com, Cell Phone:+923234682639

## ABSTRACT

**Objective:** To determine the prevalence of deficiency of vitamin D levels in first trimester of pregnancy among primigravida women.

**Study Design:** Prospective, Cross-sectional

**Place and Duration:** Conducted in the department of Obstetrics & Gynaecology, Amna Inayat Medical College / Kishwar Fazal Teaching Hospital, Lahore for duration of six months from July 2020 to December 2020.

**Methods:** Total 80 primigravidas women were included in this study. Patients' detailed demographics including age, body mass index were recorded after taking written consent. Enrolled patients were aged between 18-45 years of age. Women in their first trimester of pregnancy were presented in this study. Serum 25-hydroxy vitamin D3 [25 (OH) D3] level were observed by taking blood samples from all patients. Complete data was analyzed by SPSS 22.0 version.

**Results:** Mean age of the patients were 23.8 ±4.7 years with mean BMI 20.08±2.91 kg/m<sup>2</sup>. Out of 80 primigravidas women in their first trimester of pregnancy, deficiency in serum vitamin D level (25 (OH) D3 <30ng/mL) was found among 72 (90%) patients and only 8 (10%) women were found sufficient serum vitamin D level. Among 72 deficient women, frequency of severely deficient (<10ng/ml) women were 11(13.75%), deficient women (10-<20ng/ml) were 15(18.75%) and insufficient women (20-<30ng/ml) were 46 (57.5%). Mean serum level of vitamin D deficiency was 14.1±2.34.

**Conclusion:** We concluded in this study that the prevalence of deficiency of vitamin D levels in first trimester of pregnancy among primigravida women were significantly high.

**Keywords:** First trimester, primigravida, Vitamin D, pregnancy

---

## INTRODUCTION

Vitamin D deficiency has almost been a constant epidemic in recent times. The problem does not impact only developed countries but it has also become an immense problem in developing countries.[2] If there are issues that are so common among men and women, why do we not have a pregnant concern of population? It's a problem of about a milliard of people worldwide, according to an estimate. Especially in Pakistan, where we do not understand the state of the art or specific preventive or surveillance plan during pregnancy.

Vitamin D has its place in calcium metabolism and is a fat soluble vitamin. It is metabolised by the specially measured liver into 25-hydroxy vitamin D3 [25(OH) D3]. Collectively, the serum calculation reflects vitamin D achieved from exposure to sun and diet. Serum level is required at 30ng/ml (75nM/L), but the bone issue can only be prevented at 20ng/ml.[3] These normal parameters should be alerted to a non-pregnant population, and the normal pregnancy conditions must also be established .[4]

The function of vitamin D has to do not only with the integrity of bones, but with preeclampsia, pregnancy and other adverse results[5], it has also been found that in the mother hypovitaminosis D of the newborns, low vitamin D levels have been observed in the cord blood. The long-term neonatal storehouses of Vitamin D also rely on their foetal levels. Thus we stop hypovitaminosis in the neonate at birth and childhood by controlling the mother's vitamin D. For the maternal preservation, as well as for the formation

and development of foetal skeletons and chronic disease after birth and in later life, short vitamin D is required. [6]

Serum vitamin D levels in general depend on your diet, sun exposure, and skin content of clothing and melanin. However, is sun exposure and the normal 200 IU intake adequate to satisfy increasing pregnancy requirements? In recent decades[8] studies have been centered both in Western countries and in countries like Saudi Arabia, Iran and India, in which women wear covered clothes. The causes of the deficiency can variable, however, from place to place[7]. The figures were higher than anticipated. The conditions worsened in winter, particularly when outdoor activities were scarce, Christopher Naugler, JianguoZhanget al [9] found.

Our women in Pakistan are little known to us. The culture of men and women must be fully clothed, and the heads, hands and feet are regularly uncovered. Women have built knowledge in Pakistani urban environments to use sun blocks and to prevent direct sunlight exposure. These interventions help prevent tanning of the skin as the beauty mark of our society is a fair skin. With regard to pregnancy, there are a minimum of vitamin D in the multi-vitamin supplements used during pregnancy (5-12,5 micrograms/d). In addition, vitamin D is not fortified in routine food, flour or dairy products. Other qualitative factors may be a regional factor, education level, exercise levels, parity, socio-economic status, smoking, maternal BMI and gestation era.[10]

## MATERIAL AND METHODS

This prospective cross-sectional study was conducted at department of Obstetrics & Gynaecology, Amna Inayat Medical College / Kishwar Fazal Teaching Hospital, Lahore for duration of six months from July 2020 to December 2020 and comprised of 80 patients. Patients detailed demographics age, body mass index, was recorded after taking written consent. Patients with medical illness, diabetes, hypertension, tuberculosis cardiac illness and those were not agreed were excluded from this study.

Enrolled patients were aged between 18-45 years of age. Women in their first trimester of pregnancy were presented in this study. Serum 25-hydroxy vitamin D3 [25 (OH) D3] level were observed by taking blood samples from all patients. LMP measured gestational age and obstetric ultrasonography confirmed it. Information was registered, such as age, educational level, gestational age, parity, income monthly, smoking, exposure daily to the sun or use of multivitamins. Complete data was analyzed by SPSS 22.0 version.

## RESULTS

Mean age of the patients were 23.8  $\pm$ 4.7 years with mean BMI 20.08 $\pm$ 2.91 kg/m<sup>2</sup>. Out of 80 primigravida women in their first trimester of pregnancy, deficiency in serum vitamin D level (25 (OH) D3 <30ng/mL) was found among 72 (90%) patients and only 8 (10%) women were found sufficient serum vitamin D level. (table 1)

Table 1: Baseline detailed demographics of enrolled cases

Variables	Frequency (n=80)	%age
Mean age	23.8 $\pm$ 4.7	
mean BMI	20.08 $\pm$ 2.91	
Vitamin D Deficiency	72	90
No Deficiency	8	10

Among 72 cases of deficient pregnant women, frequency of severely deficient (<10ng/ml) women were 11(13.75%), deficient women (10-<20ng/ml) were 15(18.75%) and insufficient women (20-<30ng/ml) were 46 (57.5%). Mean serum level of vitamin D deficiency was 14.1 $\pm$ 2.34. (table 2)

Table 2: Distribution of cases with respect to vitamin D serum level

Variables	Frequency (n=80)	%age
Mean serum level	14.1 $\pm$ 2.34	
<b>25(OH)D levels</b>		
Sufficient (30ng/ml)	8	10
Insufficient (20-<30ng/ml)	46	57.5
Mild deficient(10-<20ng/ml)	15	18.75
Severe deficient(<10ng/ml)	11	13.75
<b>Total</b>	80	

## DISCUSSION

Deficiency in vitamin D during pregnancy is very common. Our study found that the prevalence of vitamin D deficiency was (90 %) comparable to that observed in Khurshid N et al and IPehlivan et al, [11,12] while South Carolina study reported 41 %of pregnant females with deficiency of vitamin D[13] The study also found that in 62%Caucasian pregnant women and96%in African American pregnant women had deficiencies in vitamin D [13,14]

A study indicates a statistically significant correlation between women with vitamin D deficiency and housewives (90 percent), who had insufficient sunlight exposure (85 %), no vitamin D deficiency supplementation (90 %), aged between 18-45 years, living in urban areas (81 %) and multiparous women (70% ).These results go with a study done by Dipali P et al [15]. The risk of vitamin D deficiency in housewives who tended to live indoors was higher and direct sunlight exposure was decreased. The shortcoming is that the exposure based on sunlight still has a bias for religion. The burqa practise is more vulnerable to vitamin D deficiency in the Muslim culture. An significant factor affecting the status of vitamin D is likely to be sunlight exposure at different latitudes. The results of previous studies of vitamin D status among pregnant women in Asia at variable latitudes and in other regions near 36°N, [16-18]which is similar to the latitude of the this study.

In our research, the fact that educated working women were less likely to have vitamin D deficiency compared to uneducated women or housewives was taken into account by socio-demographic factors. It has had its own impact on lifestyle and socioeconomic status. Pregnancy itself requires higher vitamin D levels. And all pregnant women appear to be at risk for vitamin D deficiency.[19]The deficiency is usually due to insufficient food consumption or inadequate exposure to the sun. There is no screening protocol for the amount of vitamin D in standard treatment during the prenatal period. No wonder the studies of South Asia's sun-enriched countries also have remarkable data during pregnancy on vitamin D deficiency.[20] Because knowledge and prompt response can easily adjust the risk factors, pregnant women can be encouraged to carefully monitor their intake of vitamin D and to make easy changes to their lifestyle.

The distinctive feature of our research is that with no known medical disease or concurrent drug treatment, we enrolled young, healthy primigravida. But the drawback of our studies is that we have not observed the full duration of pregnancy, which might have helped to predict levels of cord blood in delivery. In recent years, the possible effect of vitamin D deficiency during pregnancy on maternal and neonatal health has attracted a lot of interest. However, using Hill's criteria, a causal relation between vitamin D deficiency during pregnancy and adverse pregnancy-related results remains to be determined.

## CONCLUSION

We concluded in this study that the prevalence of deficiency of vitamin D levels in first trimester of pregnancy among primigravida women were significantly high.

## REFERENCES

1. FoudaMA, Turkestani IZ, Almusharraf S, Al-Ajlan A, Angkaya-Bagayawa FF, Sabico S, Mohammed AG, Hassanato R, Al-Serehi A, Alshingetti NM, Al-Daghri NM. Extremely high prevalence of maternal and neonatal vitamin D deficiency in the Arab population. *Neonatology*. 2017;112(3):225-30.
2. Saraf R, Morton SM, Camargo Jr CA, Grant CC. Global summary of maternal and newborn vitamin D status—a systematic review. *Maternal & child nutrition*. 2016 Oct;12(4):647-68.

3. El Hayek Fares J, Weiler HA. Implications of the nutrition transition for vitamin D intake and status in Aboriginal groups in the Canadian Arctic. *Nutrition reviews*. 2016 Sep 1;74(9):571-83
4. McCarthy EK, Kiely M. Vitamin D and muscle strength throughout the life course: a review of epidemiological and intervention studies. *Journal of human nutrition and dietetics*. 2015 Dec;28(6):636-45
5. Dawodu A, Davidson B, Woo JG, Peng YM, Ruiz-Palacios GM, Guerrero MD, Morrow AL. Sun exposure and vitamin D supplementation in relation to vitamin D status of breastfeeding mothers and infants in the global exploration of human milk study. *Nutrients*. 2015 Feb 5;7(2):1081-93.
6. Galthen-Sørensen M, Andersen LB, Sperling L, Christesen HT. Maternal 25-hydroxyvitamin D level and fetal bone growth assessed by ultrasound: a systematic review. *Ultrasound in Obstetrics & Gynecology*. 2014 Dec 1;44(6):633-40.
7. Fares S, Sethom MM, Khouaja-Mokrani C, Jabnoun S, Feki M, Kaabachi N. Vitamin A, E, and D deficiencies in tunisian very low birth weight neonates: prevalence and risk factors. *Pediatrics & Neonatology*. 2014 Jun 1;55(3):196-201
8. Hilger J, Friedel A, Herr R, Rausch T, Roos F, Wahl DA, Pierroz DD, Weber P, Hoffmann K. A systematic review of vitamin D status in populations worldwide. *British Journal of Nutrition*. 2014 Jan;111(1):23-45.
9. Marshall I, Mehta R, Petrova A. Vitamin D in the maternal-fetal-neonatal interface: clinical implications and requirements for supplementation. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2013 May 1;26(7):633-8
10. Naugler C, Zhang J, Henne D, Woods P, Hemmelgarn BR. Association of vitamin D status with socio-demographic factors in Calgary, Alberta: an ecological study using Census Canada data. *BMC public health*. 2013 Dec;13(1):316.
11. Khurshid N, Rafique A, Nisar F. Prevalence of Hypovitaminosis D Among Primigravidas in First Trimester of Pregnancy. *J. Soc. Obstet. Gynaecol. Pak*. 2018; Vol 8(3):159-163.
12. Pehlivan I, Hatun S, Aydoğan M, Babaoglu K, Turker G, Gökalp AS. Maternal vitamin D levels in the third trimester of pregnancy. *Turk J med Science*. 2002;32:237-241.
13. Bodnar LM, Simhan HN, Powers RW, Frank MP, Cooperstein E. High prevalence of vitamin D insufficiency in black and white pregnant women residing in the northern united states and their neonates. *J Nutr*. 2007;137(2):447-452.
14. Bowyer L, Catling-Paull C, Diamond T, Homer C, Davis G, Craig ME. Vitamin D, PTH, and calcium levels in pregnant women and their neonates. *Clin Endocrinol (Oxf)*. 2009;70(3):372-377
15. Prasad D, Smita, Singh K, Nisha S. Vitamin D in pregnancy and its correlation with fetomaternal outcome. *International Journal of Contemporary Medical Research*. 2018;5(1):1-5.
16. Fernandez-Alonso A.M., Dionis-Sanchez E.C., Gonzalez-Salmeron M.D., Perez-Lopez F.R., Chedraui P. First-trimester maternal serum 25-hydroxyvitamin D(3) status and pregnancy outcome. *Int. J. Gynaecol. Obstet*. 2012;116:6-9. doi: 10.1016/j.ijgo.2011.07.029.
17. Perez-Lopez F.R., Fernandez-Alonso A.M., Ferrando-Marco P., Gonzalez-Salmeron M.D., Dionis-Sanchez E.C., Fiol-Ruiz G., Chedraui P. First trimester serum 25-hydroxyvitamin D status and factors related to lower levels in gravids living in the Spanish Mediterranean coast. *Reprod. Sci*. 2011;18:730-736. doi: 10.1177/1933719110396720.
18. Ustuner I., Keskin H.L., Tas E.E., Neselioglu S., Sengul O., Avsar A.F. Maternal serum 25(OH) D levels in the third trimester of pregnancy during the winter season. *J. Matern. Fetal Neonatal Med*. 2011;24:1421-1426. doi: 10.3109/14767058.2011.566768.
19. Saadiya A. Karim, Uzma Nusrat and Sina Aziz. Vitamin D deficiency in pregnant women and their newborns as seen at a tertiary-care center in Karachi, Pakistan. *International Journal of Gynecology & Obstetrics*, 112, 1, (59-62), (2010).
20. Ojah RC, Welch JM. Vitamin D and musculoskeletal status in Nova Scotian women who wear concealing clothing. *Nutrients*. 2012 May 24;4(5):399-412.