

Compare the Outcomes of Sclerotherapy with Bleomycin Versus Surgical Resection in Children with Cystic Hygroma

MUHAMMAD DARAZ KHAN¹, FARIN ZADA², MASOOD SHAH³, FAROOQ ABDULLAH⁴, SAFDAR ALI⁵, HAYAT UR REHMAN⁶

¹Assistant Professor of Paediatric Surgery Bannu Medical College Bannu/ Khalifa Gul Nawaz Medical Teaching Institute, Bannu

²District Specialist Paediatric Surgery, DHQ hospital, Buner

³Consultant Paediatric Surgeon DHQ Hospital, Khar Bajure

⁴Assistant Professor Paediatric Surgery Lady Reading Hospital, Peshawar

⁵Medical Officer Paediatric Surgery, DHQ hospital, Buner

⁶Associate Professor of Paediatric Surgery Nowshera Medicare College, Nowshera

Corresponding Author: Dr Muhammad Daraz Khan, Email: pedssurgery71@yahoo.com, Cell No: 03312862628

ABSTRACT

Objective: To compare the outcomes of sclerotherapy with bleomycin versus surgical resection in children with cystic hygroma.

Study Design: Comparative/Observational

Place and Duration: Paediatric department of Khalifa Gul Nawaz Medical Teaching Institute, Bannu for 20 months duration from 1st August 2019 to 31st March 2021.

Methods: Total 40 patients of both genders with ages 1 month to 15 years presented with cystic hygroma were included in this study. Patients' detailed demographics were recorded after written consent from parents/guardians. Patients were divided in to two groups' i.e. Group A consist of 20 patients and received sclerotherapy with intra-lesional bleomycin and Group B consist of 20 patients received surgical excision. Outcomes were analyzed at post-procedure 3 and 6 months post-procedure and compared the findings between both groups.

Results: There was no significant difference observed between both groups regarding age, gender and site of lymphangioma p-value >0.05. In Group A 15 (75%) patients showed excellent results, 3 (15%) patients showed good results and 2 (10%) showed poor results. In Group B 13 (65%) showed excellent results, 4 (20%) showed good results and 3 (15%) showed poor results. At final follow-up there were no patient with recurrence in Group A while in Group B 2 (10%) patients had recurrence,

Conclusion: It is concluded that sclerosing treatment with bleomycin is a safe and effective treatment modality with no recurrence as compared to surgical resection.

Keywords: Cystic Hygroma, Sclerotherapy, Bleomycin, Surgery, Recurrence

INTRODUCTION

Lymphangiomas are benign hamartomatouslymphatic tumors, also referred to as congenital deformities of the lymphatic structures[1]. The most common and frequent type of lymphangiomas is cystic hygroma which may compose of one or more macrocytic lesions with reduced communication to the lymphatic channels [1]. They are slow growing tumors and may manifest in any part of the body or anywhere in the soft tissues.

The most commonly affected sites are the head and neck, and also the mediastinum and axilla[2]. These tumors most often occur in children, although they may occur in adults as well. They are also often evident at birth in 65% and may present by two years of age in 80-90% of the cases. The reported incidence of lymphangiomas is 1.5 to 2.8 per 1000, and it is not specific to any gender or race[2]. Inflammation and cosmetic defects are the baseline symptoms. In the neck area, they may lead to compression symptoms when they compress important surrounding structures and may cause respiratory problems, dysphagia, and compression of nerves[3].

There are different modes of treatment for cystic hygromas. These therapies include radiotherapy, incision and drainage, and surgical excision (which is the mainstay of therapy) [2,4-6]. However, these procedures have variable findings which investigators find unsatisfactory. In many patients, a complete surgical

resection is often not possible because of the nature of the lesion which encircles the crucial neurovascular system and also infiltrates across tissue planes[2]. Post-surgical recurrence of tumors and injury to nerves are commonly observed complications, in addition to other issues such as disfigurement, and dreadful scar formation [1,2].

In recent times, intralesional sclerotherapy has become increasingly accepted as an effective method for the treatment of the children with lymphangiomas [1, 2, 4]. It involves the use of a sclerosing agent which irritates the endothelial lining of the lymphangioma leading to swelling, involution, and fibrosis[2]. In the past, boiling water, 50% dextrose water, hypertonic saline, or absolute alcohol have been used with results that have not been very encouraging[2]. Many centers are beginning to use sclerosing agents like Bleomycin, acetic acid, OK-432, and Doxycycline as first-line therapy with satisfactory results [1,3,7,8]. Unlike surgical excision, this modality of treatment is particularly useful for lesions enveloping vital structures [8]. The present study was conducted to compare the effectiveness and safety of sclerotherapy and surgical excision in children with cystic hygroma.

MATERIAL AND METHODS

This observational study was conducted at Paediatric department of Khalifa Gul Nawaz Medical Teaching Institute, Bannu for 20 months duration from 1st August

2019 to 31st March 2021. A total 40 patients of both genders with ages 1 month to 15 years presented with cystic hygroma were included in this study. Patients detailed demographic including age, sex, and site of lymphangioma were recorded after taking written consent from parents/guardians. Patients with recurrence, already on sclerotherapy and ages above 15 years were excluded from this study.

All the patients were divided in to two groups' i.e. Group A consisted of 20 patients and received sclerotherapy intra-lesionally bleomycin with a dose of 0.3mg/kg of body weight and number of sessions 1 to 4 per patient. Patients were admitted for 24 hours after each session. Maximum 4 sessions were given to patients monthly. Group B consisted of 20 patients who received surgical excision of cystic hygroma. Post-procedural complications were recorded. Outcomes in terms of excellent (Complete resolution), good (>50% resolution) and poor (<50% resolution) results were examined. Recurrence rate was examined at final follow-up. Follow-up was taken at 3 and 6 months post-procedure. Ultrasonography and X-ray was done pre and post-operatively to analyze the outcomes. Data was analyzed by SPSS 24.0. Chi-square and student t test was used to compare the outcomes between both groups. P-value <0.05 was considered as statistically significant.

RESULTS

There were 11 (55%) male and 9 (45%) were females in Group A and in Group B 12 (60%) patients were males and 8 (40%) patients were females. In group A majority 75% of patients were ages <5 years and 25% patients had ages above 5 years. In Group B 80% patients were ages < 5 years and 20% patients had ages above 5 years. According to the site of lymphangioma, in Group A 60% of patients had a neck. 15% had axilla, 15% had face and 10% had trunk site and in Group B 55% patients had neck, 20% had axilla, 15% had face and 10% patients had trunk lymphangioma. There was no significant difference observed regarding age, sex and site of lymphangioma between both groups with p-value >0.05. (Table 1)

Table No 1. Baseline details of all the patients

Characteristics	Group A (n=20)	Group B (n=20)	P-value
Gender			>0.05
Male	11 (55)	12 (60)	
Female	9 (45)	8 (40)	
Age			>0.05
<5 years	15 (75)	16 (80)	
>5 years	5 (25)	4 (20)	
Site of Lymphangioma			>0.05
Neck	12 (60)	11 (55)	
Axilla	3 (15)	4 (20)	
Face	3 (15)	3 (15)	
Trunk	2 (10)	2 (10)	

In Group A patients 1 (5%) patients had received twosessions of bleomycin, 4 (20%) had received threesessions and 15 (75%) had received 4 doses of bleomycin. (Table 2)

According to the post-procedural complications we found no patient hadwound infection in Group A patients

while in Group B 2 (10%) patients had developed wound infection and 2 (10%) patients had recurrence. There was a significant difference between both groups. (p=<0.001)

Table No 2. Sessions wise distribution in patients receive bleomycin

Variables	Frequency No.	%age
One Session	0	0
Two Sessions	1	5
Three Sessions	4	20
Four Sessions	15	75

Table No 3. Post-operative complications between both groups

Characteristics	Group A (n=20)	Group B (n=20)	P-value
Wound Infection			>0.001
Yes	0 (0)	2 (10)	
No	20 (100)	18 (90)	
recurrence			>0.001
Yes	0 (0)	2 (10)	
No	20 (100)	18 (90)	

According to the resolution we found no significant difference between both groups in terms of complete resolution with p-value >0.05. In Group A 15 (75%) patients showed excellent results, 3 (15%) patients showed good results and 2 (10%) showed poor results. In Group B 13 (65%) showed excellent results, 4 (20%) showed good results and 3 (15%) showed poor results. (Table 4)

Table No 4. Final outcomes between both groups

Variables	Group A (n=20)	Group B (n=20)	P-value
Excellent	15 (75)	13 (65)	>0.05
Good	3 (15)	4 (20)	>0.05
Poor	2 (10)	3 (15)	>0.05

DISCUSSION

Cystic hygroma is one of the critical disorders among children and the incidence rate is quite high in children with ages up to 5 years. Many treatment modalities such as sclerotherapy with bleomycin and OK-432 and the surgical excision have been applied for this benign disorder. In these modalities sclerotherapy technique is considered as much safer and effective than surgical management due to high rate of wound infection and recurrence rate [9-10]. The present study was conducted aimed to examine the outcomes of sclerotherapy with bleomycin and surgical excision in children with cystic hygroma. In this regard 40 patients of both genders were enrolled and divided into two groups. We found that male patients were high in numbers in both groups A and B 55% and 60% as compared to females 45% and 40%. In group A majority 75% of patients were ages <5 years and 25% patients had ages above 5 years. In Group B 80% patients were ages < 5 years and 20% patients had ages above 5 years. A study conducted by Mustafa G et al [11] regarding outcomes of intralesional bleomycin for cystic hygroma in children, in which they reported that male patients was high in numbers 66.7% as compared to females and the mean age of patients was 2.36 ± 2.8 years.

In present study according to the site of lymphangioma, in Group A 60% patients had neck. 15% had axilla, 15% had face and 10% had trunk site and in

Group B 55% patients had neck, 20% had axilla, 15% had face and 10% patients had trunk lymphangioma. There was no significant difference observed regarding age, sex and site of lymphangioma between both groups with p-value >0.05. A study by Fiaz M et al [12] reported that neck was the commonest site of lymphangioma found in 63.3% followed by axilla, face and trunk 13.3%, 13.3% and 10%.

In our study we found that of patients who received sclerotherapy, 1 (5%) patients had received two sessions of bleomycin, 4 (20%) had received three sessions and 15 (75%) had received 4 doses of bleomycin. These results were similar to many other studies in which mostly patients received 3 to 4 sessions of bleomycin for complete resolution [13-14].

In present study According to the post-procedural complications we found no patient had wound infection in Group A patients while in Group B 2 (10%) patients had developed wound infection and 2 (10%) patients had recurrence. There was a significant difference between both groups

($p < 0.001$). These results were similar to many of previous studies in which surgical excision had high rate of wound infection 5 to 20% and recurrence rate 10 to 30% as compared to sclerotherapy [15-16]. We observed that surgical excision needs much per-operative care as compared to sclerotherapy.

In this study, according to the resolution we found no significant difference between both groups in term of complete resolution with p-value >0.05. In Group A 15 (75%) patients showed excellent results, 3 (15%) patients showed good results and 2 (10%) showed poor results. In Group B 13 (65%) showed excellent results, 4 (20%) showed good results and 3 (15%) showed poor results. These results were comparable to several previous studies [17-18].

CONCLUSION

We concluded from this study that sclerosing treatment with bleomycin for cystic hygroma in children is a safe and effective treatment modality with no recurrence and wound infection as compared to surgical resection. Also we didn't observe any significant difference between both groups in terms of complete resolution, good and poor resolution.

REFERENCES

1. Kumar V, Kumar P, Pandey A, Gupta DK, Shukla RC, Sharma SP, et al. Intralesional, bleomycin in lymphangioma: an effective, and safe non-operative modality of treatment. *J CutanAesthet Surg*. 2012;5: 133-6.
2. Mirza B, Ijaz L, Saleem M, Sharif M, Shaikh A. Cystic hygroma: an overview. *J CutanAesthet Surg*. 2010;3:139-44.
3. Rozman Z, Thambidorai RR, Zaleha AM, ZZakaria, ZulfiqarMA. Lymphangioma: is intralesional bleomycin sclerotherapy effective? *Biomed Imaging Interv J*. 2011;7:18.

4. Manikoth P, Mangalore G P, Megha V. Axillary Cystic Hygroma. *J Postgrad Med* 2004;50:215-6.
5. Okada A, Kubota A, Fukuzawa M, Imura K, Kamata S. Injection of bleomycin as a primary therapy for cystic lymphangioma. *J Pediatr Surg*. 1992 Apr;27(4):440-3.
6. Sanlialp I, Karnak I, Tanyel FC, Senocak ME, Buyukpamukcu N. Sclerotherapy for lymphangioma in children. *Int J Pediatr Otorhinolaryngol*. 2003;67:795-800.
7. Erikçi V, Hoşgör M, Yıldız M, Örnek Y, Aksoy N, Okur Ö, Demircan Y, Genişol İ. Intralesional bleomycin sclerotherapy in childhoodlymphangioma. *Turk J Pediatr*. 2013 Jul-Aug;55(4):396-400.
8. Yetiser S, Karaman K. Treatment of lymphangioma of the face with intralesional bleomycin: case discussion and literature review. *J Maxillofac Oral Surg*. 2011;10:152-4.
9. World Medical Association. World Medical Association Declaration of Helsinki Ethical Principles for Medical Research Involving Human Subjects. *JAMA*. 2013;310(20):2191-2194.
10. Redkar RG, Chigicherla S, Joshi S, Bangar A, Tewari S. Efficacy of intralesional bleomycin as an alternative approach in the management of vascular anomalies. *Saudi Surg J* 2017;5:60-4.
11. Mustafa G, Mirza B, Ahmad S, Talat N, Saleem M, Sheikh A. Outcome of Intra-Lesional Bleomycin in the Management of Cystic Hygroma. *Pak Pediatr J* 2018; 42(4): 261-65.
12. Fiaz M, Bshir A, Irfan Ullah. Effectiveness of Intralesional Bleomycin Sclerotherapy for Lymphangioma in Children. *JMSCR Volume 06 Issue 01 January 2018*.
13. Saha AK, Haque SS, Islam KM. Effect of intralesional bleomycin as an alternative therapy for cystic hygroma. *Bangladesh Med J Khulna* 2013;46:12-5.
14. Jain V, Mohta A, Sengar M, Bansal K, Valvi K. Use of intralesional bleomycin as primary therapy in macrocysticlymphangiomas. *Indian J DermatolVenereolLeprol* 2013;79:524-5.
15. Abdur-Rahman LO, Awolaran O, Nasir AA, Bamigbola KT, Abdulraheem NT, Oyinye AO, Adeniran JO. Efficacy of bleomycin for non-operative treatment of cervical lymphangioma in University of Ilorin Teaching Hospital, Nigeria. *J Med Trop* 2017;19:93-7.
16. Hassan H, Aly KA. Management of cystic lymphangioma: experience of two referral centers. *Annals of Pediatric Surgery*. 2012; 8:123-128.
17. Alqahtani A, Nguyen LT, Flageole H, Shaw K, Laberge JM. 25 years' experience with lymphangiomas in children. *J Pediatr Surg*. 1999;34(7):1164-8.
18. Bilci S, Avedi V, Oztas T, Asena M and Celik K. The Treatment of Lymphangiomas with Bleomycin in Childhood: A Retrospective Observational Study. *HaydarpasaNumune Med J* 2019;59(4):315-319.
20. Chen YN, Chen CP, Lin CJ, Chen SW. Prenatal Ultrasound Evaluation and Outcome of Pregnancy with Fetal Cystic Hygromas and Lymphangiomas. *J Med Ultrasound* 2017;25:12-5.
21. Bagrodia N, Defnet AM, Kandel JJ. Management of lymphatic malformations. *CurrOpinpediatr* 2015;27:356-63.