

# Frequency of Complications After ECT in Patients with Schizophrenia

HARESH KUMAR<sup>1</sup>, MAHIRA SHAFI<sup>2</sup>, RAKESH KUMAR<sup>3</sup>, OM PARKASH<sup>4</sup>, SARIKA BAI<sup>5</sup>, AZRA SHAHEEN<sup>6</sup>

<sup>1</sup>Assistant Professor, Psychiatry, Khairpur Medical College, Khairpur Mirs

<sup>2</sup>Assistant Professor, Institute of Psychiatry, Baqai Medical University, Karachi

<sup>3</sup>Senior Registrar, Psychiatry, Hamdard Medical University, Karachi.

<sup>4</sup>Liaquat University of Medical and Health Sciences, Jamshoro

<sup>5</sup>Woman Medical Officer, Tamachani Maternity Home, Sukkur

<sup>6</sup>Assistant Professor, Clinical Psychologist, Baqai Medical University Karachi

Corresponding Author: Dr. Haresh Kumar, Email: hareshmakhija@gmail.com, Mob: 03337589901

## ABSTRACT

**Objective:** The aim of this study is to determine the frequency of complications after ECT in patients with schizophrenia.

**Study Design:** Retrospective study

**Place and Duration:** Institute of Psychiatry Baqai Medical University, Karachi for sixteen months duration from June 2019 to September 2020.

**Methods:** Total 120 patients of both genders were presented in this study. Patients were aged between 20-70 years. Patients detailed demographics age, sex and mean BMI were recorded after taking informed written consent. Patients of schizophrenia received electroconvulsive treatment (ECT). Frequency of immediate complications was observed after each session of ECT and at the end of ECT frequency of long term complications were observed. Complete data was analyzed by SPSS 24.0 version.

**Results:** Total 65 (54.17%) were males and 55 (45.83%) were females in this study. Mean age of the patients were 40.14±3.45 years with mean BMI 22.14±6.12 kg/m<sup>2</sup>. Mean ECT sessions was 88.13±6.87. Mean hospitalization stay was 3.4±2.04 weeks. Frequency of immediate complications were 25 (20.83%) among patients after ECT session. Among 20.83%, frequency of body aches was 7(8.83%), headache was in 11 (9.17%), frequency of transient amnesia was among 3 (2.5%) and hypertension was among 4 (3.33%). Significantly no any delay complications were observed in continuously sessions among enrolled cases.

**Conclusion:** We concluded in this study that the use of ECT sessions in patients of schizophrenia was effective because no any delay complications were observed in this treatment.

**Keywords:** Schizophrenia, ECT, Treatment Sessions

---

## INTRODUCTION

Introduced in 1938 and used for almost 80 years, electroconvulsive therapy (ECT). It requires an electrical current in the brain through the head, which causes an asthma seizure and 1 muscle relaxant. ECT is widely used in various severe, therapeutically refractory or treating-resistant psychiatric conditions, including large depressive 2, schizophrenia, and bipolar disorders, as one of the most important biological therapies. Research has shown that ECT has caused major changes to the physiology and chemistry of the molecular brain, accounting for its 3 therapeutic impact.

A retrospective graphic examination of 19,982 inpatients aged between 18 and 59 years found that ECT was 66.3% in severe depression, 55.2% in schizophrenia, 68.4% in bipolar disorders and 28.6% in other psychiatric disorders[4]. China is the most frequent patient with ECT. A longitudinal research conducted in 1,364 years of age and older staff showed that in schizophrenia, the share of ECT was 57.0 percent, in severe depression 53.4 percent, in bipolar disorder 57.8 percent and in other diagnoses 32.4 percent[5]. Zhang et al. investigated ECT prevalence in Chinese adolescent psychiatric patients, and concluded that for schizophrenia, the incidence of ECT use was 46.5%, for bipolar, 41.8% and 23.9% for other diagnoses[6].

In adult patients with schizophrenia, electroconvulsive therapy is safe and effective particularly when a quick response is needed[7]. Some ECT effectiveness trials showed important advantages in the treatment of

schizophrenia in children and adolescents [8,9]. Current neurosciences show that while issues with short-term memory also arise after electroconvulsive therapy (ECT), there is a reduced prevalence of long-term problems. However, we do have shortcomings with regard to long-term cognitive issues following ECT, like memory[10]. The key drawback of the wider use of ECT is cognitive effects, especially occasional acute confusion shortly post-treatment, retrograde amnesia and some long-term autobiographic (personal) memory losses[11]. Some memory elements are either unchanged or enhanced.

ECT is practiced markedly differently from more advanced countries in countries with reduced resources; while ECT administration without anesthesia is not ideal, it is often appropriate to apply it in view of severe financial constraints. The ECT administration analysis was carried out on 1520 patients (29% of total facility admissions, 1352 (88.9%) of whom had no kind of anesthesia and no adverse reactions[12] had been registered during the procedure). Using the Clinically Global Impression-Severity (CGI-S) questionnaire the severity of each patient's psychopathology before and after ECT was measured. The aggression levels before and after the therapy were assessed using the Scale-Revised Staff Observation Aggression (SOAS-R)[13] and the operation levels before and after therapy were assessed using a Global Function Assessment (GAF) questionnaire. The treating doctors also stated the reasons for the selection of several therapies for ECT in these patients and their care schemes. These questionnaires were completed.

## MATERIAL AND METHODS

This retrospective study was conducted at Institute of Psychiatry Baqai Medical University, Karachi for sixteen months duration from June 2019 to September 2020 and comprised of 120 patients. Baseline details of patients including age, sex and body mass index (BMI) were calculated after taking written consent. Patients who had chronic medical illness (cardiovascular, respiratory disease) and those did not give written consent were excluded from this study.

Patients were selected for ECT when they showed no improvement on pharmacotherapy alone. Patients of schizophrenia were aged between 20-70 years. Frequency of immediate complications was observed after each session of ECT and at the end of ECT frequency of long term complications were observed. Categorical variables were assessed by frequency and percentage but descriptive variables were calculated by standard deviation. Complete data was analyzed by SPSS 24.0 version.

## RESULTS

Total 65 (54.17%) were males and 55 (45.83%) were females in this study. Mean age of the patients were 40.14±3.45 years with mean BMI 22.14±6.12 kg/m<sup>2</sup>. Mean ECT sessions was 88.13±6.87. Mean hospitalization stay was 3.4±2.04 weeks. (table 1)

Table 1: Characteristics of enrolled cases

Variables	Frequency	%age
<b>Sex</b>		
Male	65	54.17
Female	55	45.83
Mean Age	40.14±3.45	-
Mean BMI	22.14±6.12	-
Mean Hospitalization	88.13±6.87	-
Mean ECT Sessions	3.4±2.04	-

Among 120 patients, most of the patients 66.7% were from the age group 20-40 years and received higher number of ECT sessions while remaining 33.3% were >40 years and received minimum sessions. (table 2)

Frequency of immediate complications were 25 (20.83%) among patients after ECT session. Among 20.83%, frequency of body aches was 7(8.83%), headache was in 11 (9.17%), frequency of transient amnesia was among 3 (2.5%) and hypertension was among 4 (3.33%). Significantly no any delay complications were observed in continuously sessions among enrolled cases. (table 3)

Table 2: Association of ECT with respect to age

Variables	Frequency (n=120)	% age
<b>Age group</b>		
20-40 years (maximum sessions)	80	66.7
>40 years (minimum sessions)	40	33.3

Table 3: Frequency of complications among schizophrenia patients

Variables	Frequency (n=120)	% age
<b>Complications</b>	<b>25</b>	<b>20.83</b>
Body Aches	7	8.83
Headache	11	9.17
Transient amnesia	3	2.5
Hypertension	4	3.33

## DISCUSSION

Though ECT is well known and evidence-based in the acute phase of major psychiatric conditions, ECT knowledge is lacking. In this retrospective study, complications were collected between 120 schizophrenia patients. Total 65 (54.17%) were males and 55 (45.83%) were females in this study. Mean age of the patients were 40.14±3.45 years with mean BMI 22.14±6.12 kg/m<sup>2</sup>. Mean ECT sessions was 88.13±6.87. Mean hospitalization stay was 3.4±2.04 weeks. These findings were comparable to the many previous studies. [14,17] While Lévy-Rueff et al.[18] reported M-ECT in schizophrenic or schizo-affective patients with a 48±20 year mean age, Suzuki et al.[17] reported M-ECT in a collection of schizophrenia with 61 year medium age patients.

Our study only included schizophrenic patients. There were no patients with bipolar affective disorder who received multiple ECT in our hospital. This is either due to the fact that patients who have affective distress typically do not have to have a lot of ECT, the fact that doctors treating them do not provide multiple ECTs, or because of the technical and logistical complexities involved in offering multiple ECT treatments. This research covers a special patient group and the literature contains few reports on these patients. A group of schizophrenia or schizoaffective disorder patients who had been hospitalized an average 10 months before M-ECT were identified by Lévy-Rueff et al [18] but, on average, hospitalized for two months after initial care. Lévy-Rueff et al. Russell et al[19] reported having been in the group most of the year before beginning C/M-ECT their series of bipolar affective and schizoaffective patients.

In our study frequency of immediate complications were 25 (20.83%) among patients after ECT session. Among 20.83%, frequency of body aches was 7(8.83%), headache was in 11 (9.17%), frequency of transient amnesia was among 3 (2.5%) and hypertension was among 4 (3.33%). Significantly no any delay complications were observed in continuously sessions among enrolled cases. These were comparable to the previous studies. [20-23] While ECT under anesthesia is ideal, it is hard to do in any patient even in countries such as Pakistan, where there is a lack of resources. There seems to be no distinction between the two ECT modalities in delayed complications. Furthermore, ECT is often determined for patients with extreme mental conditions that can place suicidal patients, stupor, catatonia or serious self-neglect of their own or others. Therefore ECT will save lives in such a situation. [24,25]

The side-effects and complications observed in other studies[17,19] are common, but are minor when found, including in elderly and sometimes physically ill patients. Minor side effects, such as headaches, nausea and muscles, are no different from those recorded after acute ECT treatment. Keys, long apnea, heart dysrhythmias, epileptic or delirium status, and other major complications are very uncommon. Both of these need an expert team to closely track each ECT treatment and any one after it.

ECT can cause some adverse effects, including memory impairment and new learning, long-term seizures, general anesthesia risk and other minor effects. Some reports, however, do not agree with this view. De Serna et

al. found no substantial improvement discrepancies between the ECT community and the non-ECT group over a two-year follow-up in clinical or neuropsychological variables over time. There is also no detrimental effect of ECT in their study on long-term neuropsychological variables. [26] The report [26] Lima et al. found that ECT is highly effective for treating multiple adolescent psychological conditions with low and relatively mild effects in a systematic review of 39 trials[27]. Further studies on the adverse effects of ECT, particularly in young people, are needed.

## CONCLUSION

We concluded in this study that the use of ECT sessions in patients with schizophrenia was effective because no any delay complications were observed in this treatment.

## REFERENCE

1. Saleem R, Rauf S and Jehangir S. Efficacy of Electroconvulsive Therapy in treatment resistant Depression. *Pak Armed Forces Med J* 2018; 68 (4): 969-74.
2. Leiknes KA, Jarosh-von Schweder L, Hoie B. Contemporary use and practice of electroconvulsive therapy worldwide. *Brain Behav.* 2012;2(3):283–344.
3. Singh A, Kar SK. How Electroconvulsive Therapy Works?: Understanding the Neurobiological Mechanisms. *Clin Psychopharmacol Neuro Sci* 2017; 15(3): 210-21.
4. Wang ZM, Zhu H, Pan YL, Chiu HF, Correll CU, Ungvari GS, Lai KY, Cao XL, Li Y, Zhong BL, Zhang XY, Xiang YT. Electroconvulsive therapy and its association with demographic and clinical characteristics in Chinese psychiatric patients. *J ECT.* 2015;31(2):114–8.
5. An FR, Zhang L, Zhang QE, Ungvari GS, Ng CH, Chiu HFK, Wu PP, Jin X, Li L, Lok GKI, Xiang YT. Electroconvulsive therapy and its relationships with clinical characteristics and quality of life in Chinese psychiatric patients. *Psychiatry Res.* 2016;246:246–9.
6. Zhang QE, Wang ZM, Sha S, Ng CH, Seiner SJ, Welch CA, Lok GK, Chow IH, Wang F, Li L, Xiang YT. Common use of electroconvulsive therapy for chinese adolescent psychiatric patients. *J ECT.* 2016;32(4):251–5.
7. Tharyan P, Adams CE. Electroconvulsive therapy for schizophrenia. *Cochrane Database Syst Rev.* 2005;18:CD000076.
8. Baeza I, Flamarique I, Garrido JM, et al. Clinical experience using electroconvulsive therapy in adolescents with schizophrenia spectrum disorders. *J Child AdolescPsychopharmacol.* 2010;20:205–9.
9. Maoz H, Nitzan U, Goldwyn Y, Krieger I, Bloch Y. When can we predict the outcome of an electroconvulsive therapy course in adolescents? A Retrospective Study. *J ECT.* 2018;34:104–7.
10. Coetzer R. Depression, memory and electroconvulsive therapy. *BJPsych Bull.* 2019 Apr; 43(2): 51–53.
11. Semkowska, M. and D. M. McLoughlin (2010). "Objective cognitive performance associated with electroconvulsive therapy for depression: a systematic review and meta-analysis." *Biol Psychiatry* 68(6): 568-577.
12. Minhas Mh ,OstroffR.Practice of electroconvulsive therapy in a tertiary care hospital in Pakistan *J ECT.* 2012 Mar;28(1):7-9.
13. Nijman H, Palmstierna T. Measuring aggression with the Staff Observation Aggression Scale-Revised. *Acta PsychiatrScand Suppl.* 2002;412:101–102.
14. Iancu I, Pick N, Seener-Lorsh O, Dannon P. Patients with schizophrenia or schizoaffective disorder who receive multiple electroconvulsive therapy sessions: characteristics, indications, and results. *Neuropsychiatr Dis Treat.* 2015;11:853-862. Published 2015 Mar 27. doi:10.2147/NDT.S78919
15. Wang, S., Yang, C., Jia, J. et al. Use of electroconvulsive therapy in adolescents with schizophrenia in China. *Child Adolesc Psychiatry Ment Health* 12, 49 (2018).
16. O'Connor DW, Gardner B, Presnell I, Singh D, Tsanglis M, White E. The effectiveness of continuation-maintenance ECT in reducing depressed older patients' hospital re-admissions. *J Affect Disord.* 2010;120:62–66.
17. Suzuki K, Takano T, Ebina Y, Takamatsu K, Awata S, Matsuoka H. Continuation electroconvulsive therapy to prevent relapse of schizophrenia in relapse-prone patients. *J ECT.* 2007;23:204–205.
18. Lévy-Rueff M, Gourevitch R, Loo H, Olié JP, Amado I. Maintenance electroconvulsive therapy: an alternative treatment for refractory schizophrenia and schizoaffective disorders. *Psychiatry Res.* 2010;175:280–283.
19. Russell JC, Rasmussen KG, O'Connor M, Copeman CA, Ryan DA, Rummans TA. Long-term maintenance ECT: a retrospective review of efficacy and cognitive outcome. *J ECT.* 2003;19:4–9.
20. UK ECT Review group Efficacy and safety of electroconvulsive therapy in depressive disorders: a systematic review and metaanalysis *Lancet*2003 Mar 8;361(9360):799-808.
21. Coetzer R. Depression, memory and electroconvulsive therapy. *BJPsych Bull.* 2019 Apr; 43(2): 51–53.
22. Semkowska, M. and D. M. McLoughlin (2010). "Objective cognitive performance associated with electroconvulsive therapy for depression: a systematic review and meta-analysis." *Biol Psychiatry* 68(6): 568-577.
23. Brown SK, Nowlin RB, Sartorelli R., Smith J, Johnson K. Patient experience of electroconvulsive therapy: a retrospective review of clinical outcomes and satisfaction. *J ECT* 2018, PMID: 29613943.
24. ECT Handbook (3rd Edition). 2013 Edited by Waite, J. & Easton, A. 2013. The Royal College of Psychiatrists.
25. Pagnin, D., V. de Queiroz, S. Pini and G. B. Cassano (2004). "Efficacy of ECT in depression: a meta-analytic review." *J ECT* 20(1): 13-20.
26. De la Serna E, Flamarique I, Castro-Fornieles J, Pons A, Puig O, Andrés-Perpiña S, Lázaro L, Garrido JM, Bernardo M, Baeza I. Two-year follow-up of cognitive functions in schizophrenia spectrum disorders of adolescent patients treated with electroconvulsive therapy. *J Child AdolescPsychopharmacol.* 2011;21(6):611–9.
27. Lima NN, Nascimento VB, Peixoto JA, et al. Electroconvulsivetherapy use in adolescents: a systematic review. *Ann Gen Psychiatry.* 2013;12:17.