

ORIGINAL ARTICLE

Dissimilar Perspectives and Facts about Procedure of Amenorrhea Between Fledgling Ladies

HASEEB UL HASSAN¹, KHIZRA MANZOOR², QURATULAIN HAIDER³, TAHIRA FATIMA⁴, MOINUDDIN AHMED⁵, NAUMAN AHMED⁶

^{1,2}*Surgery Department, Abbas Institute of Medical Sciences Muzaffarabad Azad Kashmir*

³*Surgery Department, Abbas Institute of Medical Sciences, Muzaffarabad*

⁴*Associate Professor, Gynae and Obs Department, Shalamar Medical and Dental College, HOD Fauji foundation Hospital Lahore*

⁵*HIESS, Hamdard University, Karachi*

⁶*PGR Eye Unit 1, Lahore General Hospital, Lahore*

Correspondence to: Haseeb Ul Hassan, Email: haseeb.0079@gmail.com, Cell: 03469077771

ABSTRACT

Objective: This investigation work aimed to find out the awareness and numerous perspectives towards the procedure of mensuration between ladies of fledgling age.

Methodology: This descriptive investigation work carried out in Jinnah Hospital, Lahore and its remote areas from December 2017 to November 2018. The selection of the 500 ladies carried out from remote areas of hospital and students of this institute with the help of non-probability sampling method. Ladies present with the abnormalities in amenorrhea procedure or suffering from other gynecological problems were excluded from this investigation work. All the selected ladies were in their menstruating age regardless of their marriage and education stage. SPSS V. 22 was in use for the arithmetical exploration of the collected information. We recorded all the information of a well-organized Performa.

Results: Out of total five hundred participants, 84.5% (n: 415) ladies stated that this procedure is beneficial for health; however, 18.5% (n: 85) felt that this incident was not healthy. Out of total five hundred ladies, 88.65% (n: 438) ladies regarded the procedure of amenorrhea as normal procedure, while 13.45% (n: 62) ladies thought this as God's curse.

Conclusion: Health education of the ladies particularly living in the rural areas is much necessary. Health education will prepare the ladies emotionally to face this procedure of amenorrhea with less adverse reactions. Faulty perception, un-factsment and unsecure practices about the procedure of amenorrhea are very common between ladies of fledgling age.

Keywords: Arithmetical, exploration, unsecure, Sampling, amenorrhea, Performa, abnormalities.

INTRODUCTION

This procedure describes the onset and end of the potential of reproductive system [1]. The phenomenon of amenorrhea is unique to women [2]. Perceptions and compartments towards procedure of amenorrhea may present with variations in dissimilar cultures [3]. Amenorrhea is cyclic endometrium's shedding [4] and it also plays a vital part in the concepts of menstrual distress [5]. Specifically speaking, the cycles of amenorrhea are the indicators of the health of ladies [6]. Amenorrhea on monthly basis is natural, beneficial and normal for ladies [7, 8], Perspectives and compartments towards the procedure of amenorrhea may badly impact the body image of ladies, concept of the causation of disease, dietary habits, consent to get medication, usage of contraception and ability to have pregnancy planning. [9]. amenorrhea is not a procedure of discrete incidence but it is a procedure of cycle of events in the whole life time.

Discussing the procedure of amenorrhea with the peers was associated to grade to which this procedure is thought to be a procedure of nature and the sharing of the experience of amenorrhea will enhance the perception of these procedure as natural [10]. The procedure of amenorrhea is the distinct sign of potential of reproduction and maturity of sexual life. We stated that the perspectives of the ladies towards the procedure of amenorrhea will have correlation with the sexual compartment, compartment and destinies [11]. The proper comprehension of clinical and cultural features about the

amenorrhea in our populace is not much common and talking about this procedure is considered as taboo. One of the most important issue in Muslims ladies of our country Pakistan is that because of the social and economic humiliation, the ladies neither wants to reveal their health issues nor they visit gynecologist and they normally experience issues like high loss of blood causing anemia of severe nature [12]. The restriction to unlock the cultural as well as psychosocial taboos is a predominant confusion. This leads to ignorance of the amenorrhea as a natural procedure and it can lead to difficulties in managing such issues. Woman may feel about self-conscious in the duration of their amenorrhea procedure because they normally view this procedure as social humiliation [13].

In populaces of our communities, a host of blend messages about this natural procedure of amenorrhea are provided to the pre-pubertal fledgling girls. The available data about this procedure is contradictory because it congratulates the fledgling ladies on their entrance in to the womanhood whereas at the same duration, it is suggested that this should be kept confidential from others [14]. The main objective of this investigation work was to determine the compartments and facts of ladies about the amenorrhea between ladies of fledgling age and to give the providers of health care field with basic strategies to educated the ladies and support them in making healthy and hygienic choices. [15]. In comparison with the societies of west, our fledgling girls do not have much education well in time for these vital alterations in their lives and all these

ignorant fledgling ladies can give reaction in a complete dissimilar way

METHODOLOGY

This investigation work carried out in Jinnah hospital, Lahore on students of the institute and the ladies of remote regions. This investigation work carried out on fledgling ladies of society. Total five hundred ladies in the menstruating age regardless of their education stage and marriage status got recruitment for managing a well-organized questionnaire in this investigation work. This investigation work started in December 2018 to November 2019. All the ladies present with abnormalities in the procedure of amenorrhea or suffering from some other mental issues and gynecological anomalies got exclusion from this investigation work. We used non-probability sampling method for the selection of the participants.

SPSS V. 22 was in use for the arithmetical exploration of the collected information. We presented the age in average and standard deviations. We calculated all the quantitative variables in percentages and frequencies. We collected all the information of the samples on well-organized Performa, information included data about age of the female, marriage status, profession, and stage of education as well as facts about the procedure of amenorrhea and practices regarding this procedure.

RESULTS

The range of the age of the ladies was from fourteen to thirty eight years. Total five hundred fledgling ladies were participants of this investigation work. The average age of the ladies was 23.956 ± 5.030 years. Monthly income of fifty percent participants was greater than ten thousand rupees. We performed the descriptive exploration of the social and demographic variables to describe the traits of our populace. Total 49.85% (n: 247) ladies were from urban regions whereas remaining ladies were from non-urban areas of the city. Only 5.25% ladies were working while all other ladies were present with no occupation. The average age at the time of menarche was 12.825 ± 1.155 years. There were total 52.45% (n: 263) participants were married whereas remaining participants were unmarried as elaborated in (Table-1).

Table 1: Socio demographic data

Variables	No. of Victims	Percentage (%)
Age (Years)	>30	10.4
	< 20	50.5
	21-30	40.6
Monthly household income (PKR)	Upper (>10,000)	50.7
	Poor (<5000)	40.5
Educational Status	Middle (5000-10,000)	9.5
	Metric	0.7
	FA/FSC	1.8
	Graduate	36.9
	Nil	59.8
	Primary	1.8
Middle	0.6	





Table-2 displays the menstrual traits of the studied populace. The lowest age at the time of menarche was nine years and the highest age at the time of menarche was sixteen years.

Table 2: Menstrual Characteristics

Information Regarding Amenorrhea	Participants	Percentage
Awareness From	Teacher	5 (1.02)
	Literature	5 (1.05)
	Others	32 (6.22)
	No information	116 (23.25)
	Mother	276 (55.22)
Perception of Amenorrhea	Others (Disease & curse)	62 (12.42)
	Natural procedure	438 (87.65)
Is Amenorrhea necessary for women?	Yes	393 (78.65)
	No	107 (21.45)
Opinion About Cessation of Amenorrhea with Drugs	Yes	20 (4.02)
	No	480 (96.05)
Amenorrhea Good For Health	Yes	415 (83.05)
	No	85 (17.05)



About 77.40% (n: 387) ladies stated that they do not take some particular food in this period whereas 22.60% (n: 113) stated that they used to take hot soups, vegetables

and fruits. Most of the ladies 53.25% (n: 268) were embarrassed at the time of menarche. Total 38.25% (n: 191) ladies stated that they preferred to avoid sexual activities, 16.44% (n: 82) ladies avoided bath, 17.7% (n: 85) ladies avoided to take part in physical activity in the period of amenorrhea. They stated the extent of the impact as mil, mild, moderate and severe as 49.66% (n: 248), 33.65% (n: 168), 14.66% (n: 73), and 2.24% (n: 11) correspondingly. We asked the Victims about the impact of this amenorrhea on the physical activities.

DISCUSSION

Showed the same results in which emotional reactions of the participant toward menarche were hugely negative and about 85.50% ladies were annoyed and embarrassed. About the perception of the amenorrhea procedure, 87.60% (n: 438) ladies thought it to be a procedure of nature for cleaning but Mc Master in his investigation work found that only 12.5% ladies consider this procedure as a cleaning system [16]. This current investigation work stated that there are some adverse practices, false perceptions and misconceptions in ladies of fledgling age and these same issues have been stated by many other investigation works conducted in numerous countries of the world. Majority of the ladies 53.25% (n: 266) felt embarrassed as a response to the menarche whereas in a investigation work conducted by Tang CS [17]. 17.5% (n: 85) ladies avoided to participate in physical activity, whereas investigation work conducted by Snow LF [18] stated that 37.55% ladies thought that cold air and water should be avoided in the duration of amenorrhea procedure. We discovered that numerous practices which mostly ladies avoid in the procedure of amenorrhea include avoiding sexual activities by majority of ladies 38.25% (n: 191), 16.45% (n: 82) ladies avoided to take bath.

CONCLUSION

The findings of this investigation work concluded that health education about this procedure requires to be stated at early ages before the puberty age of the ladies. Health education also reduces the negative reactions of the fledgling ladies. This health education will also make them able to better care in the procedure of amenorrhea.

REFERENCES

1. Nelson LM, Hillard P, Trott J, Khosla S, Warren MP. The menstrual cycle-getting the "vital" respect it deserves. [2003]

- [cited 15 Feb, 2008]. Available from: http://www.pituitary.org/news/the_menstrual_cycle_getting_the_vital_respect.aspx.
2. McPherson ME, Korfine L. Amenorrhea across time: menarche, menstrual perspectives, experiences, and compartments. *Womens Health Issues* 2004;14:193-200.
 3. Rempel JK, Baumgartner B. The relationship between perspectives towards amenorrhea and sexual perspectives, desires, and compartment in women. *Arch Sex Behav* 2003;32:155-63.
 4. Anson O. Exploring the bio-socil approach to premenstrual experiences. *Soc Sci Med* 1999;49:67-80.
 5. Kowalski RM, Chapple T. The social humiliation of amenorrhea. *Psychol Women Q* 2000;24:7480.
 6. Perveen F, Hashmi H. Dysfunctional uterine bleeding: A clinic histopathological exploration based management. *J Coll Physicians Surg Pak* 1999;9:18-20.
 7. Dan AJ. What have we learned? An historical view of the society for menstrual cycle investigation. *NWSA J* 2004;16:45-55.
 8. Beausang CC, Razor AG. Fledgling western women's experiences of menarche and amenorrhea. *Health Care Women Int* 2000;21:517-28.
 9. Tang CS, Yeung DY, Lee AM. Psychological correlates of emotional responses to menarche between Chinese adolescent girls. *J Adolesc Health* 2003;33:193-201.
 10. McMaster J, Cormie K, Pitts M. Menstrual and premenstrual experiences of women in a developing country. *Health Care Women Int* 1997; 18:533-41.
 11. Snow LF, Johnson SM. Modern day menstrual folklore: Some clinical implication. *J Am Med Assoc* 1977;237:2736-9.
 12. Drakshayani Devi K, Venkata Ramaiah P. A study on menstrual hygiene between rural adolescent girls. *Indian J Med Sci* 1994;48:139-43.
 13. Kubba A. Amenorrhea, cultural perspectives and medical aspects: An over view. *Gynaecol Forum* 2003;8:3-6.
 14. Chohan A. Puberty, amenorrhea, and ovulation. In: *Fundamentals of Gynaecology*. Lahore MAR Publications 2000:15-20.
 15. Lu ZJ. The relationship between menstrual perspectives and menstrual symptoms between Taiwanese women. *J Adv Nurs* 2001;33:621-8.
 16. Chandra PS, Chaturvedi SK. Cultural variations in perspectives towards amenorrhea. *Can J Psychiatry* 1992;37:196-198.
 17. Thoms SL, Ellertson C. Nuisance or natural and healthy: should monthly amenorrhea be optional for women? *Lancet* 2000;335:922-4.
 18. Fort IL, Brezzo RD. Menstrual cycle alteration with aging. *Am J Health Studies* 1999;15:41-6.