

# Effect of Trunk Stabilization Exercises on Static and Dynamic Sitting Balance among Children with Cerebral Palsy: A Randomized Control Trial

FAIQA MEHMOOD RANA<sup>1</sup>, SAIMA JABBAR<sup>2</sup>, AMNA ZIA<sup>3</sup>, RABBIA AKHTAR<sup>4</sup>, MAHAM KHALID<sup>5</sup>, SANA UMAR<sup>6</sup>

<sup>1</sup>School of Health Sciences, Department of Rehabilitation Medicine, University of Management and Technology, Pakistan

<sup>2</sup>The Green International University, Lahore, Pakistan.

<sup>3</sup>School of Physiotherapy, King Edward Medical University, Mayo Hospital, Pakistan

<sup>4</sup>Department of Physiotherapy, University of South Asia, Pakistan

<sup>5</sup>School of Health Sciences, Department of Rehabilitation Medicine, University of Management and Technology, Pakistan

<sup>6</sup>University of health sciences

Correspondence to Dr. Faiqa Mehmood Rana, Email: [faiqa.rana1996@gmail.com](mailto:faiqa.rana1996@gmail.com)

## ABSTRACT

**Aim:** To evaluate the effectiveness of trunk stabilization exercises on Static and Dynamic Sitting Balance among Children with Cerebral Palsy

**Method:** This randomized controlled trial was conducted at physiotherapy department, Mayo hospital, Lahore. 38 children with cerebral palsy were enrolled in this study via non-probability purposive sampling technique which later on randomized and allocated to two equal groups. Group A received specific core stability training while group B received conventional physical therapy treatment. Written informed consent was procured from every participant or their guardian through signed consent forms. Questionnaire used for data collection was Gross Motor Function Measure (GMFM-88) also used for assessment of sitting balance.

**Results:** Descriptive statistics showed participants were having mean age of  $6.00 \pm 2.00$  for group A and  $5.52 \pm 1.80$  for Group B. Participants who received trunk stabilization exercises showed marked improvement as compared to group B. There was a statistically significant improvement in all measured variables following the intervention ( $P < 0.05$ ). Pre-treatment sitting GMFM mean score in group A was  $27.54 \pm 3.61$ , while in group B was  $26.48 \pm 4.64$ . After 3 week treatment mean score in group A was  $30.04 \pm 3.81$  and in group B was  $27.43 \pm 4.36$  and post treatment mean score in group A was  $32.80 \pm 3.54$  and in group B was  $29.46 \pm 4.36$ .

**Practical Implication:** This research would lay another evidence highlighting the effectiveness of trunk stability exercises to not only improve sitting balance among cerebral palsy patients but also the quality of life among them.

**Conclusion:** It was concluded from the study that trunk stabilization exercises showed significant improvement in static and dynamic sitting balance scale among cerebral palsy children. The children when treated with trunk stabilization exercises showed a significant treatment outcome when analyzed on Gross Motor Function Measure GMFM-88.

**Keywords:** Cerebral Palsy, Trunk control, Static sitting Balance, Dynamic sitting balance, GMFM-88.

## INTRODUCTION

Cerebral Palsy includes permanent neurological disorders that lead to motor, sensory, cognitive issues and ultimately activity limitations. Such neurological problems are associated with non-progressive abnormalities or lesions occurred in early stages of brain development<sup>1</sup>. Optimal trunk stability is considered vital to help execute basic functional activities of daily life like sitting and reaching. Impaired trunk control is a characteristic hallmark which often exists among patients of cerebral palsy but there is deficient evidence about the extent of impairment in static and dynamic trunk stability in moderate to severe cases of cerebral palsy<sup>2</sup>.

The trunk's insufficient function is a very important factor that render difficulties not only in basic activities of daily living like sitting, playing and gross movements but also in performing complex functional daily living tasks such as moving in a wheelchair<sup>3</sup>. Voluntary control of the trunk usually used to represent an important stage in development of function and for some children it may leads to independent activity in sitting, so that independent sitting balance without specialized seating may be achieved<sup>4</sup>. The trunk muscle strengthening exercise is considered effective to improve the functional gross movement and the balance in children having spastic cerebral palsy<sup>5</sup>. The management and improvement of the core muscles strength, coordination and balance have also enhanced overall efficacy in functions of kinetic chains of upper and lower extremities. Reciprocally, inefficient core stability could have been considered to correspond to decreased functional capabilities affecting gross motor skills in CP child<sup>6,7</sup>. Muscle strengthening is considered very effective in improving sitting posture, gross motor function and the functional independence. Muscle strengthening has shown positive effects on the postural stability in sitting<sup>8</sup>.

Postural control is being achieved by subconscious coordinated muscle activity in space to stabilize the body's position or optimize postural alignment. Postural control usually plays a primary role during functional movements and the daily tasks<sup>9</sup>. Therefore, it is considered very necessary to evaluate the trunk control ability in these positions. Trunk sway is being correlated with the functional activities in children with cerebral palsy<sup>10</sup>.

Postural stability along with reaching movements are usually considered dependent on the extent of segmental trunk control that is acquired during sitting<sup>11</sup>. Trunk training mostly targets activation of core and lower extremity muscles increasing their strength, improve balance as well as gait problems<sup>12</sup>. The Muscle activation showed increased in the bilateral erector spine, rectus abdominal muscles in the left RA, bilateral RF, gastrocnemius, and left TA. Trunk Training is being considered safely and effectively maintains and improves physical performance and mostly can be included in the rehabilitation programs<sup>13</sup>.

Spinal sagittal alignment activation is associated with the improvement in the functional outcomes and with the reduced complications rates. That's why trunk instability had led to deforming forces on the spine and ultimately scoliosis<sup>14</sup>. Treatment usually necessitates a cautious risk-benefit analysis. Therefore it is considered imperative to have in-depth indulgence in patients' medical history and post-morbid functional status<sup>15</sup>.

Neck and trunk stabilization exercises had shown a positive impact on static sitting balance, movement control and dynamic sitting balance<sup>16</sup>.

The rationale of this study lies in the focus on trunk stabilization exercises and the development of an intervention protocol to benefit patients and to prove that trunk stabilization exercises would improve not only trunk function but also static and dynamic sitting balance, to a greater extent than seen with a conventional comprehensive rehabilitation. This study helped to justify that how strongly trunk musculature is linked with balanced movement of extremities in children with cerebral palsy.

Received on 19-06-2022

Accepted on 09-11-2022

**MATERIALS & METHODS**

Present study was chosen to be a randomized controlled trial. After approval from ethical committee of Mayo Hospital, the study procedures were conducted at school of physiotherapy, Mayo hospital. Informed consent had been taken from all participants and their caregivers through signed consent forms. Thirty-eight children with spastic paraplegic CP were selected. The patients were aged 3–10 years. There was random manual assignment of patients to either group; an intervention group or a control group.

Sample size was calculated by employing G\*POWER(version 3.1.9.4.) statistical tool. The estimated sample size was found to be 19 participants in each group. Calculations were assumed using  $\alpha=0.05$ , Effect size  $d = 1.1043153$ , power  $=0.95$ . To overcome dropout ratio, sample size was increased. Mean of group A is 0.4 and mean of group B is 0.45. Standard deviation of group 1 is 0.04 and standard deviation of group 2 is 0.05<sup>17</sup>.

Inclusive criteria for the study was Spastic CP with paraplegia and impaired trunk control. Participants aged (3 to 10) years with mild degree of spasticity (grade 1 to 1+ according to Modified Ashworth Scale), Level II or III of the Gross Motor Function Classification system GMFCS were included.

The exclusive criteria for the study were participants aged more than 10 years, having sitting position already intact, any Co morbidities affecting motor performance.

**Treatment interventions:** Performa was collected and participants were received an informed consent. Questionnaire used to record assessment data was GMFM-88 which was used for assessment of sitting balance Group A received Trunk Stabilization Exercises (Figure 1) along with conventional Physiotherapy three times/week for total six weeks and Group-B received Conventional Physiotherapy, three times/week for total six weeks treatment duration.

One group was assigned as intervention or Group A, whose participants were treated with trunk stabilization training with each single session lasting for about forty five minutes. Standard exercise protocol designed by Jeffreys<sup>18</sup>, which was incorporated in this group included three levels. First two levels included static and dynamic balance exercises in a stable environment whereas, third level comprised of dynamic and resistive exercise training amongst unstable environment as discussed in Figure 1, along with conventional physiotherapy treatment and other group was treated with only the conventional rehabilitation program. In this study hot pack was applied to all of patients in every treatment session<sup>19</sup>. General muscle stretching exercises for upper and lower limb (typically for calf muscles by pulling the toes down toward the floor and lifting the heel off the ground, other muscles like adductor muscles and muscles of the arm) provided to the patients of both groups as conventional physiotherapy treatment. The patients were treated 6 weeks duration (3 times per week) each treatment session lasted for 25 min.

Assessment of the participants were done at baseline (pretreatment), mid-treatment and post-treatment following six weeks of intervention. The assessor was taken on board who was not directly involved in treatment provision and was also kept blind to the allocation of each child to their respective group. The assessor was a practicing, qualified and an experienced pediatric physical therapist as well as an efficient administrator of GMFM. Though special training WAS not necessary to administer the tool, researches recommend that the administrator must be familiar with assessment of children motor skills and the general GMFM administration protocol<sup>20</sup>.The patients were re-evaluated by the same assessor at the end of the treatment.

**Outcome Measures:**

**GMFM-88 (Gross Motor Function Measure-88):** Outcome measure, used in this study, has been considered as being fantastic tool for approaching an initial assessment. We have used GMFCS to determine Gross Motor skills and to find how severe a given case of cerebral palsy was. This scoring system might

helped in taking consideration of limb control , movement, shifting position , use of assistive devices and many other factors<sup>21</sup>. Gross Motor Function Measure scale has been the gold standard outcome measure to assess the level of gross motor skills and functioning among patients with cerebral palsy at clinical and as well as research settings<sup>22</sup>. It has exhibited well-established and excellent psychometric properties for the population having cerebral palsy in children (age< 15 years) and has also been validated against other populations as well. As this is an assessment tool, could be effectively applied to record change in motor abilities and functions over time as well as to reflect the efficacy of interventions devised to improve gross motor skills<sup>23</sup>.

**Static and Dynamic Sitting Balance:** Static balance, when defined, is the capability of child to uphold the body at rest while being in any stable or fixed posture and to carry postural alignment in coordinated and stabilized way keeping center of mass within base of support<sup>24</sup>.On the other hand, dynamic posture has been quiet challenging for children with cerebral palsy, capability to fluctuate the center of gravity within vertical projection around line of gravity and base of support. In short, to maintain body posture with well oriented balance and coordination keeping it within base of support while it is in motion, is referred to as dynamic balance<sup>25</sup>.

**Statistical analysis:** Using SPSS 23, data analysis was executed. Demographic data was computed using mean and standard deviation to interpret the descriptive statistics. For quantitative data, mean and standard deviation were preferred execution while qualitative categorical data has been presented in the form of percentage or frequency table if applicable. Shapiro-wilk test has been used to assume normal distribution of data. After assuming normality, parametric hypothesis testing was considered and independent sample t test was used to assess mean difference between groups while within subject variance, time effect and interaction effect, repeated measure ANOVA test was used. P-value<0.05 was considered significant to reject null hypothesis.

Figure 1: Trunk Stabilization Exercises Protocol



**RESULTS**

Participants were evaluated at the time of enrollment for eligibility and baseline assessment; 46 children, fulfilling the inclusion criteria, were randomly allocated to two equal groups (Figure 2: CONSORT flow diagram). Eight patients did not comply with the study protocols due to unavoidable reasons (mentioned in Figure 2). Hence, 38 participants' data were analyzed statistically.

Table 1 shows that out of 19 (100%) subjects, 8 were male and 11 were female in group A. In Group B out of 19 (100%) subjects, 14 were male and 5 were female. mean age in group 1= 6.00 ± 2.00 and in group 2 = 5.52 ± 1.80. Distribution of cases according to socioeconomic status has showed out of 19 (100%) subjects; 14 belong to middle class and 5 belong to lower class. In Group II out of 19 (100%) subjects; 16 belong to middle class and 3 belong to lower class.

**Baseline or Pre-treatment Mean Difference:** Baseline assessment indicated that both groups, intervention and control, showed no significant difference in any of the concerned outcome variables ( $P > 0.05$ ).

**Pre-treatment, mid-assessment & post-intervention comparison in group A:** Pre-treatment sitting GMFM mean score was  $27.54 \pm 3.61$ . After 3 week treatment mean score was  $30.04 \pm 3.81$  and post treatment mean score was  $32.80 \pm 3.54$ .

**Pre-treatment, mid-assessment and post-intervention comparison in group B:** Pre-treatment sitting GMFM mean score was  $26.48 \pm 4.64$ . After 3 week treatment mean score was  $27.43 \pm 4.36$  and post treatment mean score was  $29.46 \pm 4.36$ .

**Pre-treatment, mid-assessment and post-intervention comparison for both groups:** Pre-treatment sitting GMFM mean score in group A was  $27.54 \pm 3.61$  and in group B was  $26.48 \pm 4.64$ . After 3 week treatment mean score in group A was  $30.04 \pm 3.81$  and in group B was  $27.43 \pm 4.36$  and post treatment mean score in group A was  $32.80 \pm 3.54$  and in group B was  $29.46 \pm 4.36$  as shown in Table 2. Hence, supported with mean difference over time indicated a significant improvement in static and dynamic sitting balance post intervention ( $P < 0.05$ ).

Table 1: Sociodemographic Characteristics of Participants

Age of patients (Years)		
Treatment Group	Mean	6.00
	Std. Deviation	2.00
	Minimum	3.0
	Maximum	9.0
Control Group	Mean	5.52
	Std. Deviation	1.80
	Minimum	3.0
	Maximum	9.0
Gender		
Female		22(57.89%)
Male		16(42.10%)
Socioeconomic status		
Lower class		8(21.01%)
Middle class		30(78.94%)

Shapiro wilk test of normality has shown that p-value for most of the variables in group 1 and 2 were more than 0.05, showing that data was normally distributed (Table 3). Thus, between group analysis, independent T, P value for pretreatment or baseline  $p = 0.44$  (non-significant) showing variables were similar at baseline. Mid treatment  $p$  value = 0.57 (non-significant). Post Treatment  $p =$

Table 5.1: Tests of Within-Subjects Effects

GMFM	Type III Sum of Squares	df	Mean Square	F	Sig.
Sphericity Assumed	325.568	2	162.784	263.906	.000
Greenhouse-Geisser	325.568	1.615	201.647	263.906	.000
Huynh-Feldt	325.568	1.726	188.615	263.906	.000
Lower-bound	325.568	1.000	325.568	263.906	.000

Table 5.2: Tests of Within-Subjects Contrasts

Source	GMFM	Type III Sum of Squares	df	Mean Square	F	Sig.
GMFM	Linear	322.761	1	322.761	355.174	.000
	Quadratic	2.807	1	2.807	8.641	.006

Table 5.3 Tests of Between-Subjects Effects

Source	Type III Sum of Squares	df	Mean Square	F	Sig.
Intercept	95634.61	1	95634.61	1964.10	.000

0.014 (Significant) showing treatments A were more effective than treatment B in improving the sitting balance in CP child. As it is clear that the post-assessment results showed  $p$  value significant (Table 4)

Table 5 shows multivariate analysis interpreting repeated measure ANOVA to observe time effect, within subject and between subjects comparison, results turned out to be significant ( $p$  value  $< 0.05$ ).

Table 2: Pre and Post treatment comparison in Group A and B

Group	N	Mean	Std. Deviation	Std. Error Mean
<b>Pretreatment GMFM score</b>				
Group A	19	27.54	3.61	0.83
Group B	19	26.48	4.64	1.06
<b>After 3 week treatment GMFM score</b>				
Group A	19	30.04	3.81	0.87
Group B	19	27.43	4.36	1.00
<b>Post treatment GMFM score</b>				
Group A	19	32.80	3.54	0.81
Group B	19	29.46	4.36	1.00

Table 3: Test of Normality (as  $n < 50$ , Shapiro-wilk was used to assume normality of data)

	Sig.
Pre Treatment sitting GMFM score Group A	.412
After 3 week Treatment sitting GMFM Group A	.835
Post treatment sitting GMFM score Group A	.545
Pre Treatment sitting GMFM score in Group B	.516
After 3 weeks sitting GMFM score in Group B	.341
Post Treatment sitting GMFM score in Group B	.821


Table 4: Independent Samples Test

	Independent t-test		
	t	Df	Sig. (2tailed)
Pre Treatment sitting GMFM score	78	36	.44
After 3 week sitting GMFM score	1.96	36	.05
Post treatment sitting GMFM score	2.58	36	.01

Table 5: Multivariate Test

GMFM Effect	Value	F	Sig.
Pillai's Trace	.908	172.71	.000
Wilks' Lambda	.092	172.71	.000
Hotelling's Trace	9.869	172.71	.000
Roy's Largest Root	9.869	172.71	.000

Figure 2: Time &amp; Interaction effect (Mixed Factorial Analysis)



## DISCUSSION

In this randomized controlled trial, two groups of patients each group having 19 patients. One group of patients was treated with trunk stabilization training along with conventional physiotherapy treatment and other group was treated with only the conventional rehabilitation program<sup>26</sup>. General muscle stretching exercises for upper and lower limb (typically for calf muscles by pulling the toes down toward the floor and lifting the heel off the ground, other muscles like adductor muscles and muscles of the arm) provided to the patients of both groups as conventional physiotherapy treatment. The patients were treated 6 weeks duration (3 times per week) each treatment session lasted for 25 min. The progress of patients was assessed at the end of treatment. There was notable improvement observed in outcome variable, as of both static and dynamic balance during sitting posture following the intervention protocol. The static and dynamic sitting balance was assessed by GMFM-88 (Gross Motor Function Measure) scale<sup>27</sup>.

Group A of this study who received trunk stabilization training, mean pre-treatment sitting GMFM mean score was  $27.54 \pm 3.61$ . After 3 week treatment mean score was  $30.04 \pm 3.81$  and post treatment mean score was  $32.80 \pm 3.54$ .

Group B who received conventional physiotherapy treatment, mean Pre-treatment sitting GMFM mean score was  $26.48 \pm 4.64$ . After 3 week treatment mean score was  $27.43 \pm 4.36$  and post treatment mean score was  $29.46 \pm 4.36$ . The significantly notable improvement being gained in Gross motor function measure in participants of intervention group at the end of treatment regimen which had been entirely designed considering neurodevelopmental therapy principles aiming to facilitate the normal patterns<sup>28</sup>. Moreover, the post treatment results also showed remarkable improvement in patients in the control group, being comparable with the results of study of Dodd et al<sup>29</sup>.

Improvement of gross motor function activities and balance have been supported by the findings of the study by Davis & Verheyden et al which concluded that the stability of trunk is a significant core element of balance and coordination, thus helps expedite to accomplish advance motor skills. Another research by Mudie et al concluded that kinesthetic and proprioceptive training of trunk in the patient with cerebral palsy had produced better weight distribution and balance<sup>29, 30</sup>. The ultimate enhancement of trunk control happens as stabilization training is primarily comprised of certain trunk movements aimed to cause the strengthening of trunk musculature<sup>31</sup>. The remarkable positivity in post intervention results in experimental group is due to marked improvement in abdominal and back muscle strength as well as proprioception of trunk as the treatment protocol majorly involved specific activities and training, engaging the core muscles in a

static and perturbed environments<sup>32, 33</sup>. This study helped to find that the CP children with good trunk control seemed more active in static and dynamic balanced sitting. Thus it stated the importance to determine the trunk control required for the improvement of sitting balance both in static and dynamic perspective in children with cerebral palsy. Children with insufficient trunk control failed to achieve effective sitting balance that affected the functional independence.

## CONCLUSION

Post assessment of patients on GMFM-88 scale indicated that the use of trunk stability exercise regimen has produced effective and better outcomes than the routine physiotherapy program alone. The comparison of results on data analysis in my study showed that the trunk stabilization exercises along with conventional physical therapy rehabilitation program is effective in Cerebral Palsy.

**Limitations:** The major limitation of this study was poor compliance by the participants and it took longer than expected to complete the estimated sample size. Moreover, this study compared one intervention with the conventional therapy; results would have been different if compared with some other advance techniques. Sample size was not enough to produce generalizability of the results.

**Acknowledgment:** I am very thankful to the Department (School of Physiotherapy, Mayo Hospital) for their encouraging support.

**Conflict of interest:** Nil

## REFERENCES

1. Akbas AN, Gunel MK. Effects of Trunk Training on Trunk, Upper and Lower Limb Motor Functions in Children with Spastic Cerebral Palsy: A Stratified Randomized Controlled Trial. *Konuralp Tıp Dergisi*. 2019;11(2):253-9. <https://doi.org/10.18521/ktd.453532>
2. Barbado D, Reina R, Roldan A, McCulloch K, Campayo-Piernas M, Vera-Garcia FJ. How much trunk control is affected in adults with moderate to severe cerebral palsy? *Journal of biomechanics*. 2019;82:368-74. <https://doi.org/10.1016/j.jbiomech.2018.11.009>
3. Shin JW, Song GB, Ko J. The effects of neck and trunk stabilization exercises on cerebral palsy children's static and dynamic trunk balance: case series. *Journal of physical therapy science*. 2017 Apr;29(4):771-4. <https://doi.org/10.1589/jpts.29.771>
4. Butler PB. A preliminary report on the effectiveness of trunk targeting in achieving independent sitting balance in children with cerebral palsy. *Clin Rehabil*. 1998 Aug;12(4):281-93. <https://doi.org/10.1191/026921598667577442>
5. El Shemy SA. Trunk endurance and gait changes after core stability training in children with hemiplegic cerebral palsy: A randomized controlled trial. *Journal of back and musculoskeletal rehabilitation*. 2018(Preprint):1-9. DOI: 10.3233/BMR-181123
6. Kim DH, An DH, Yoo WG. Changes in trunk sway and impairment during sitting and standing in children with cerebral palsy. *Technology and health care : official journal of the European Society for Engineering and Medicine*. 2018;26(5):761-8. DOI: 10.3233/THC-181301
7. Stability and increased walking speed alters trunk movements in children with cerebral palsy. *Frontiers in human neuroscience*, 10, 354.
8. Ilharreborde, B., de Saint Etienne, A., Presedo, A., & Simon, A.-L. (2020). Spinal Sagittal alignment and head control in patients with cerebral palsy. *Journal of Children's Orthopaedics*, 14(1), 17-23. <https://doi.org/10.1302/2F1863-2548.14.190160>
9. Contini, B. G., Bergamini, E., Alvin, M., Di Stanislao, E., Di Rosa, G., Castelli, E., . . . Camomilla, V. (2019). A wearable gait analysis protocol to support the choice of the appropriate ankle-foot orthosis: A comparative assessment in children with Cerebral Palsy. *Clinical Biomechanics*, 70, 177-185. <https://doi.org/10.1016/j.clinbiomech.2019.08.009>
10. Milićević, M. (2019). Profile of motor abilities of children with cerebral palsy as a predictor of their functional independence in self-care and mobility. *Srpski arhiv za celokupno lekarstvo*(00), 77-77. <https://doi.org/10.2298/SARH190321077M>
11. Keller, J. W., & Van Hedel, H. J. (2017). Weight supported training of the upper extremity in children with cerebral palsy: a motor learning study. *Journal of Neuroengineering and Rehabilitation*, 14(1), 87



12. Acar, G., Altun, G. P., Yurdalan, S., & Polat, M. G. (2016). Efficacy of neurodevelopmental treatment combined with the Nintendo® Wii in patients with cerebral palsy. *Journal of physical therapy science*, 28(3), 774-780 <https://doi.org/10.1589/jpts.28.774>
13. Paraschiv-Ionescu, A., Newman, C. J., Carcreff, L., Gerber, C. N., Armand, S., & Aminian, K. (2019). Locomotion and cadence detection using a single trunk-fixed accelerometer: validity for children with cerebral palsy in daily life-like conditions. *Journal of neuroengineering and rehabilitation*, 16(1), 24
14. Panibatta, S., Kumar, V., & Narayan, A. (2017). Relationship between trunk control and balance in children with spastic cerebral palsy: a cross-sectional study. *Journal of clinical and diagnostic research: JCDR*, 11(9), YC05 <https://dx.doi.org/10.7860%2FJCDR%2F2017%2F28388.10649>
15. Schmid, S., Romkes, J., Taylor, W. R., Lorenzetti, S., & Brunner, R. (2016). Orthotic correction of lower limb function during gait does not immediately influence spinal kinematics in spastic hemiplegic cerebral palsy. *Gait & Posture*, 49, 457-462 2205
16. Bleyenheuft, Y., Ebner-Karestinos, D., Surana, B., Paradis, J., Sidiropoulos, A., Renders, A., Gordon, A. M. (2017). Intensive upper- and lower-extremity training for children with bilateral cerebral palsy: a quasirandomized trial. *Developmental Medicine & Child Neurology*, 59(6), 625-633 <https://doi.org/10.1016/j.gaitpost.2016.08.013>
17. Tramontano, M., Medici, A., Iosa, M., Chiariotti, A., Fusillo, G., Manzari, L., & Morelli, D. (2017). The effect of vestibular stimulation on motor functions of children with cerebral palsy. *Motor control*, 21(3), 299311 <https://doi.org/10.1123/mc.2015-0089>
18. El Shemy SA. Trunk endurance and gait changes after core stability training in children with hemiplegic cerebral palsy: A randomized controlled trial. *Journal of back and musculoskeletal rehabilitation*. 2018 Jan 1;31(6):1159-67.
19. Shin JW, Song GB, Ko J. The effects of neck and trunk stabilization exercises on cerebral palsy children's static and dynamic trunk balance: case series. *Journal of physical therapy science*. 2017;29(4):771-4. <https://doi.org/10.1589/jpts.29.771>
20. Mohamadi, O., Torabinezhad, F., & Ebadi, A. (2020). Psychometric properties of the Persian version of the oral motor assessment scale in children and adolescents with cerebral palsy. *International Journal of Developmental Disabilities*, 1-7. <https://doi.org/10.1080/20473869.2020.1819944>
21. Alotaibi M, Long T, Kennedy E, Bavishi S. The efficacy of GMFM-88 and GMFM-66 to detect changes in gross motor function in children with cerebral palsy (CP): a literature review. *Disability and rehabilitation*. 2014 Apr 1;36(8):617-27.
22. Pavão, S. L., Maeda, D. A., Corsi, C., Santos, M. M Costa, C. S. N. d., de Campos, A. C., & Rocha, N. A. C. F. (2019). Discriminant ability and criterion validity of the Trunk Impairment Scale for cerebral palsy. *Disability and rehabilitation*, 41(18), 2199-2205 <https://doi.org/10.1080/09638288.2018.1462410>
23. Pham, R., Mol, B. W., Gecz, J., MacLennan, A. H., MacLennan, S. C., Corbett, M. A., . . . Berry, J. G. (2020). Definition and diagnosis of cerebral palsy in genetic studies: a systematic review. *Developmental Medicine & Child Neurology*, 62(9), 1024-1030.
24. Chen, A., Dyck Holzinger, S., Oskoui, M., Shevell, M., & Registry, C. C. P. (2020). Losing a diagnosis of cerebral palsy: a comparison of variables at 2 and 5 years. *Developmental Medicine & Child Neurology*, 62(1), 83-88 <https://doi.org/10.1111/dmcn.14585>
25. Martakis, K., Stark, C., Rehberg, M., Semler, O., Duran, I., & Schoenau, E. (2020).
26. One-Minute Walk Test in Children with Cerebral Palsy GMFCS Level 1 and 2: Reference Values to Therapeutic Effects after Rehabilitation. *Developmental Neurorehabilitation*, 23(4), 201209. <https://doi.org/10.1080/17518423.2019.1625981>
27. Sarafian, A. J. (2020). *Development and Validation of an Observational Tool to Evaluate Upper Extremity Functioning or Hand-object Interaction in Children Diagnosed With Bilateral Cerebral Palsy GMFCS II, IV, and V*. Teachers College, Columbia University. <https://doi.org/10.7916/d8-7xdy-kt57>
28. Reference Centiles to Monitor the 6-minute-walk Test in Ambulant Children with Cerebral Palsy and Identification of Effects after Rehabilitation Utilizing Whole-body Vibration. *Developmental Neurorehabilitation*, 1-11. <https://doi.org/10.1080/17518423.2020.1770891>
29. Monbaliu, E., Himmelmann, K., Lin, J.-P., Ortibus, E., Bonouvrié, L., Feys, H., Dan, B. (2017). Clinical presentation and management of dyskinetic cerebral palsy. *The Lancet Neurology*, 16(9), 741-749. [https://doi.org/10.1016/S1474-4422\(17\)30252-1](https://doi.org/10.1016/S1474-4422(17)30252-1)
30. Chin, E. M., Gwynn, H. E., Robinson, S., & Hoon, A. H. (2020). Principles of Medical and Surgical Treatment of Cerebral Palsy. *Neurologic clinics*, 38(2), 397-416. <https://doi.org/10.1016/j.ncl.2020.01.009>
31. Carpenter, C., & Bass, A. (2017). The value of gait analysis in decision making about surgical treatment of cerebral palsy *Paediatric Orthopaedics* (pp. 361-367): Springer DOI: 10.1007/978-3-319-41142-2\_38
32. Koy, A., & Timmermann, L. (2017). Deep brain stimulation in cerebral palsy: challenges and opportunities. *European Journal of Paediatric Neurology*, 21(1), 118-121. <https://doi.org/10.1016/j.ejpn.2016.05.015>
33. Masson, R., Pagliano, E., & Baranello, G. (2017). Efficacy of oral pharmacological treatments in dyskinetic cerebral palsy: a systematic review. *Developmental Medicine & Child Neurology*, 59(12), 1237-1248. <https://doi.org/10.1111/dmcn.13532>