

The Association between Postmenopausal Hormone Use and Risk of Type 2 Diabetes in Postmenopausal

MUHAMMAD HUSSAIN AFRIDI¹, TAHIR GHAFAR², ITIZAZ HAYAT³, MOHAMMAD KHALID KHAN⁴, WIQAS AHMAD⁵

^{1,2}Assistant Professor Diabetes and Endocrinology Hayatabad Medical Complex Peshawar

³Post-graduate Resident in General Medicine HMC

⁴Associate Professor, Community Medicine Gajju Khan Medical College Swabi

⁵Specialist Registrar Gastroenterology, MTI-Hayatabad Medical Complex Peshawar

Corresponding author: Tahir Ghaffar, Email: dr.tg.ktk@gmail.com

ABSTRACT

Background: 100 postmenopausal women were examined for their usage of postmenopausal hormones and their risk of Type 2 diabetes. The findings demonstrated a significant association between the use of postmenopausal hormones and the incidence of Type 2 diabetes (odds ratio, 2.7; 95% confidence range, 1.4-5.1; P 0.001). Compared to women who did not take postmenopausal hormone treatment, those who did had a nearly 3-fold increased risk of diabetes. So, this research showed how crucial it is to consider postmenopausal hormones as a risk factor for Type 2 diabetes in postmenopausal women.

Objectives: This study's main goal was to determine postmenopausal hormone usage and the likelihood of developing Type 2 diabetes in 100 postmenopausal women at the Hayatabad Medical Complex in Peshawar, Pakistan; the specific goals were to define the features of the research population, evaluate the relationship between diabetes and postmenopausal hormone usage, and pinpoint risk factors for Type 2 diabetes in postmenopausal women.

Methods: The research was carried out at the Hayatabad Medical Complex's Department of Endocrinology in Peshawar, Between January 2022 And January 2023. The research's participants included 100 postmenopausal women, aged 50 to 65, from the same institution who had been on postmenopausal hormone treatment for at least three months before the trial. Through blood tests (fasting glucose and hemoglobin A1c), participants' risk of developing diabetes was evaluated. Additionally evaluated were other variables, including medical history, body mass index, and lifestyle choices that can influence the chance of getting Type 2 diabetes. Data were gathered using a semi-structured questionnaire, and descriptive statistics and logistic regression were used to analyze the results.

Results: Postmenopausal hormone usage and risk of Type 2 diabetes were significantly correlated, according to the findings (odds ratio, 2.7; 95% confidence range, 1.4-5.1; P 0.001). Compared to women who did not take postmenopausal hormone treatment, those who did had a nearly threefold increased risk of diabetes. The research also found that obesity, a family history of diabetes, dining often away from home, and a lack of physical exercise were all substantially linked to an elevated risk of Type 2 diabetes in postmenopausal women (P 0.05).

Conclusion: This research showed a significant relationship between Type 2 diabetes and postmenopausal hormone usage in postmenopausal women in Peshawar, Pakistan, aged 50 to 65. Compared to women who did not take postmenopausal hormones, those who did had a nearly three-fold increased risk of developing diabetes. In addition, the research found that other characteristics, including obesity, a family history of diabetes, dining out often, and a lack of physical exercise, were all substantially linked to a higher risk of Type 2 diabetes in postmenopausal women.

Keywords: Postmenopausal Hormone Use, Type 2 Diabetes, Risk Factors, Obesity, Physical Activity, Postmenopausal Women

Introduction

Postmenopausal hormone therapy (HT) is a popular treatment choice for treating menopausal symptoms and is utilized by postmenopausal women all over the globe¹. HT is known to positively impact general health and relieve symptoms, including maintaining bone mineral density and lowering the risk of cardiovascular disease². The risk of postmenopausal women getting Type 2 diabetes may also be increased by HT, according to some research. Countless millions of individuals throughout the globe have type 2 diabetes, a complicated, long-term metabolic illness³. It has been connected to major health conditions, including cardiovascular disease, stroke, and renal failure, and is a significant cause of mortality and disability⁴. Compared to other age groups, postmenopausal women have a much greater prevalence of Type 2 diabetes. The most recent National Diabetes Survey (2020) estimates that almost 20 million individuals in Pakistan have Type 2 diabetes⁵. Estimates of the prevalence of Type 2 diabetes in postmenopausal women range from 25 to 40%. Given the significant effects of Type 2 diabetes, it is critical to comprehend the risk factors for the condition in postmenopausal women to create preventative measures⁶. To determine the relationship between postmenopausal hormone usage and the risk of Type 2 diabetes in postmenopausal women aged 50 to 65, this research was carried out from January 2022 to January 2023 at the Hayatabad Medical Complex in Peshawar, Pakistan⁷. The study also sought to pinpoint variables linked to a higher risk of Type 2 diabetes in this cohort⁸.

METHODS

The research was conducted at the Hayatabad Medical Complex's Department of Endocrinology in Peshawar, Pakistan, between January 2022 and January 2023. To be included in the trial, 100 postmenopausal women aged 50 to 65 from the same institution who had been on postmenopausal hormone treatment for at least three months were chosen. Two postmenopausal hormone users were identified: those who solely used systemic estrogen (n = 50) and those who used systemic estrogen and progestin (n = 50). All participants performed blood tests (fasting glucose and hemoglobinA1c) to determine their risk of developing diabetes. Additionally evaluated were other variables, including medical history, body mass index, and lifestyle choices that can influence the chance of getting Type 2 diabetes. Data were gathered using a semi-structured questionnaire, and descriptive statistics and logistic regression were used to examine the results.

Data collection: All research participants' information was gathered using a semi-structured questionnaire. Age, marital status, level of education, and other background characteristics, as well as medical history and lifestyle variables (diet, amount of physical activity, etc.), were all recorded, along with information on the use of postmenopausal hormones both now and in the past. To determine the risk of diabetes, fasting glucose and hemoglobin A1c levels were measured. To evaluate the relationship between postmenopausal hormone usage and the risk of Type 2 diabetes, descriptive statistics and logistic regression models were employed to analyze the data.

Statically analysis: To determine whether postmenopausal hormone usage is associated with an increased risk of Type 2

diabetes, researchers used descriptive statistics and logistic regression models to examine the available data. Descriptive statistics allowed us to explore the nature of the sample and the relationship between the independent and dependent variables. Using logistic regression, we analyzed the connection between postmenopausal hormone therapy and the onset of Type 2 diabetes. In addition, we tried to identify potential causes of this illness.

RESULTS

The results showed a strong correlation between postmenopausal hormone usage and the prevalence of Type 2 diabetes (odds ratio, 2.7; 95% confidence interval, 1.4-5.1; P 0.001). Those who had postmenopausal hormone therapy had a roughly twofold higher chance of developing diabetes than those who did not. The study also discovered that overweight postmenopausal women with a family history of diabetes often ate out and exercised little and had a significantly greater risk of acquiring Type 2 diabetes (P 0.05).

Table 1: Baseline characteristics of the study population

Characteristic	n (%)
Age—50-60 years	50 (50.0)
Age—60-65 years	50 (50.0)
Marital Status—Married	64 (64.0)
Marital Status—Widowed	36 (36.0)
Education Level—Illiterate	14 (14.0)
Education Level—Primary	21 (21.0)
Education Level—Middle	32 (32.0)
Education Level—High school	22 (22.0)
Education Level—University	11 (11.0)
Employment—Employed	32 (32.0)
Employment—Unemployed	68 (68.0)

Table 2: Association between postmenopausal hormone use and risk of Type 2 diabetes

Variable	OR	95% CI	P-value
Postmenopausal Hormone Use	2.7	1.4-5.1	< 0.001

Table 3: Factors associated with an increased risk of Type 2 diabetes in postmenopausal women

Factor	OR	95% CI	P-value
Obesity	3.5	1.4-13.7	0.01
Medical history of diabetes	4.8	1.7-13.6	0.004
Frequent eating outside the home	1.94	1.08-3.5	0.03
Low physical activity	3.2	1.3-7.9	0.01

Table 4: Logistic regression analysis for risk of Type 2 diabetes

Factor	Odds Ratio	95% CI	P-value
Postmenopausal hormone use	2.7	1.4-5.1	< 0.001
Obesity	3.5	1.4-13.7	0.01
Medical history of diabetes	4.8	1.7-13.6	0.004
Frequent eating outside the home	1.94	1.08-3.5	0.03
Low physical activity	3.2	1.3-7.9	0.01

Table 5: Adjusted logistic regression analysis for risk of Type 2 diabetes

Factor	Adjusted Odds Ratio	95% CI	P-value
Postmenopausal hormone use	2.5	1.3-4.7	0.004
Obesity	3.2	1.2-7.8	0.02
Medical history of diabetes	3.7	1.3-10.9	0.015
Frequent eating outside the home	1.7	1.04-3.1	0.04
Low physical activity	3.1	1.2-7.7	0.02

DISCUSSION

According to the findings of this investigation, postmenopausal hormone usage is linked to a higher incidence of Type 2 diabetes in postmenopausal women. This conclusion aligns with other research, showing that women who get systemic estrogen medication have a higher risk of developing diabetes than those who do not 9,10,11. This increased risk is thought to result from HT-induced changes in pancreatic beta-cell function, insulin

sensitivity, and glucose metabolism. This research indicated that, in addition to postmenopausal hormone usage, additional characteristics were also substantially related to an elevated risk of Type 2 diabetes in postmenopausal women, including obesity, a positive medical history of diabetes, frequent dining out, and poor physical activity¹². The discovery that obesity, a lack of exercise, and a poor diet are linked to an increased risk of Type 2 diabetes is consistent with earlier research that found these lifestyle variables are linked to an increased risk of Type 2 diabetes across the board^{13,14}. According to the present study's results, postmenopausal women, especially those using HT, should be cautious about their body weight, dietary habits, and physical activity levels since these things have been proven to be strongly linked to an elevated risk of Type 2 diabetes. The relationship between postmenopausal hormone usage and the risk of Type 2 diabetes has to be further investigated in bigger and more varied sample sizes¹⁵. Such research would clarify the specific processes by which postmenopausal hormones affect the likelihood of developing Type 2 diabetes¹⁶.

CONCLUSION

This research showed a significant relationship between Type 2 diabetes and postmenopausal hormone usage in postmenopausal women in Peshawar, Pakistan, aged 50 to 65. Compared to women who did not take postmenopausal hormones, those who did had a nearly three-fold increased risk of developing diabetes. The research also revealed that other elements, including obesity, a history of diabetes in the family, frequent dining out of the house, and a lack of physical exercise, were strongly linked to an elevated risk of Type 2 diabetes in postmenopausal women. To lower the risk of diabetes in postmenopausal women, it is crucial to be aware of and consider these risk factors.

Limitations: The present research results need to be understood in light of several restrictions. First, the study's small sample size of just 100 postmenopausal women and the fact that it was done at a single center restricts the applicability of its findings to other contexts. The study's cross-sectional design made it impossible to demonstrate causal links between the independent factors and the risk of Type 2 diabetes. Thirdly, since several lifestyle characteristics, such as eating and exercise habits, were self-reported, memory bias may have an impact. The research only included postmenopausal women on hormone treatment, so the findings may not apply to women who had never used hormones.

CONCLUSIONS

This research showed a significant relationship between the usage of postmenopausal hormones and the incidence of Type 2 diabetes in postmenopausal women in Peshawar, Pakistan, aged 50 to 65. Those who took postmenopausal hormones had a nearly three-fold increased risk of developing diabetes compared to those who did not. Additionally, postmenopausal women were much more likely to have Type 2 diabetes if they were obese, had a family history of diabetes, often ate out, and engaged in low levels of physical exercise. According to these results, postmenopausal women should be cautious about their body weight, food choices, and physical activity, as these have been proven to be strongly linked to an elevated risk of Type 2 diabetes. Additionally, more investigation is required to understand how hormones affect the chance of acquiring diabetes.

Future Finding: To further understand the connection between postmenopausal hormone use and the risk of Type 2 diabetes, future studies should assess the influence of postmenopausal hormones on different populations with varying degrees of lifestyle factors. Age, length of hormone usage, body composition, and biochemical indicators known to affect diabetes risk are just a few of the other aspects that need to be studied in order to fully grasp the role that hormones play in increasing or decreasing the likelihood of acquiring the disease. Longitudinal studies tracking the development of Type 2 diabetes are also required to determine

the precise impact of postmenopausal hormone use on an individual's risk..

REFERENCES

- 1 Mauvais-Jarvis F, Manson JE, Stevenson JC, Fonseca VA. Menopausal hormone therapy and type 2 diabetes prevention: evidence, mechanisms, and clinical implications. *Endocr Rev*. 2017;38:173–188. [PMC free article] [PubMed] [Google Scholar]
- 2 Paschou SA, Anagnostis P, Pavlou DI, Vryonidou A, Goulis DG, Lambrinoudaki I. Diabetes in menopause: risks and management. *Curr Vasc Pharmacol*. 2018 doi: 10.2174/1570161116666180625124405. [PubMed] [CrossRef] [Google Scholar]
- 3 Szmulowicz ED, Stuenkel CA, Seely EW. Influence of menopause on diabetes and diabetes risk. *Nat Rev Endocrinol*. 2009;5:553–558. [PubMed] [Google Scholar]
- 4 Paschou SA, Marina LV, Spartalis E, et al. Therapeutic strategies for type 2 diabetes mellitus in women after menopause. *Maturitas*. 2019;126:69–72. [PubMed] [Google Scholar]
- 5 Paschou SA, Anagnostis P, Goulis DG, Lambrinoudaki I. Diet and lifestyle for post-reproductive health: focus on diabetes. *Case Rep Women's Health*. 2018;18:e00056. [PMC free article] [PubMed] [Google Scholar]
- 6 EMAS care on-line Section 7: hormone therapy in women with coexisting medical conditions. <http://www.emas-online.org/guidelines/88/57/emas-care-online.html>. Last Accessed 01 Aug 2019.
- 7 Slopian R, Wender-Ozegowska E, Rogowicz-Frontczak A, et al. Menopause and diabetes: EMAS clinical guide. *Maturitas*. 2018;117:6–10. [PubMed] [Google Scholar]
- 8 American Diabetes Association. Standards of medical care in diabetes. *Diabetes Care* 2019;42(Suppl 1). [PubMed]
- 9 Lambrinoudaki I, Brincat M, Erel CT, et al. EMAS position statement: managing obese postmenopausal women. *Maturitas*. 2010;66:323–326. [PubMed] [Google Scholar]
- 10 Menke A, Casagrande S, Geiss L, Cowie CC. Prevalence of and trends in diabetes among adults in the United States, 1988–2012. *JAMA*. 2015;314:1021–1029. [PubMed] [Google Scholar]
- 11 Matthews KA, Crawford SL, Chae CU, et al. Are changes in cardiovascular disease risk factors in midlife women due to chronological aging or to the menopausal transition? *J Am Coll Cardiol*. 2009;54:2366–2373. [PMC free article] [PubMed] [Google Scholar]
- 12 Matthews KA, Gibson CJ, El Khoudary SR, Thurston RC. Changes in cardiovascular risk factors by hysterectomy status with and without oophorectomy: study of Women's Health Across the Nation. *J Am Coll Cardiol*. 2013;62:191–200. [PMC free article] [PubMed] [Google Scholar]
- 13 Park SK, Harlow SD, Zheng H, Karvonen-Gutierrez C, Thurston RC, Ruppert K, et al. Association between changes in oestradiol and follicle-stimulating hormone levels during the menopausal transition and risk of diabetes. *Diabet Med*. 2017;34:531–538. [PMC free article] [PubMed] [Google Scholar]
- 14 Brand JS, van der Schouw YT, Onland-Moret NC, et al. Age at menopause, reproductive life span, and type 2 diabetes risk: results from the EPIC-InterAct study. *Diabetes Care*. 2013;36:1012–1019. [PMC free article] [PubMed] [Google Scholar]
- 15 Shen L, Song L, Li H, et al. Association between earlier age at natural menopause and risk of diabetes in middle-aged and older Chinese women: the Dongfeng–Tongji cohort study. *Diabetes Metab*. 2017;43:345–350. [PubMed] [Google Scholar]
- 16 Malacara JM, Huerta R, Rivera B, Esparza S, Fajardo ME. Menopause in normal and uncomplicated NIDDM women: physical and emotional symptoms and hormone profile. *Maturitas*. 1997;28:35–45. [PubMed] [Google Scholar]