

# Determine the Prevalence and Factors Contributing to Increase the Caesarean Section Rate amongst post-partum women at Lyari Town Karachi, Pakistan

SHAHLA PARDHAN SAMEER ALI<sup>1</sup>, MUHAMMAD YASIN<sup>2</sup>, SHAHID ALI KHAN<sup>2</sup>, SHABANA RASHEED BHATTI<sup>2</sup>, ESHRAT FATIMA<sup>2</sup>, TARIQ AZIZ<sup>3</sup>

<sup>1</sup>Research Faculty, School of Nursing, Kharadar General Hospital Karachi

<sup>2</sup>Post RN BSN, School of Nursing, Kharadar General Hospital Karachi.

<sup>3</sup>Nursing Instructor, School of Nursing, Kharadar General Hospital Karachi

Correspondence to Shahla Pardhan Sameer Ali, Email: [Shaileena123@hotmail.com](mailto:Shaileena123@hotmail.com), 92-3343039547

## ABSTRACT

**Aim:** To determine the prevalence and factors contributing to the increase in the Cesarean Section Rate among Post-Partum Women in Two Private Hospitals in Lyari Town, Karachi, Pakistan.

**Methodology:** A descriptive cross-sectional study was conducted from August to November 2022 after getting approval from the Institutional Review Board (IRB) of Kharadar General Hospital. The purposive sampling technique and self-structured questionnaires were used to achieve the sample size of 230 post-partum women with age < than 45 were included in the study whereas Women who were delivered vaginally, and those who were not willing were excluded in this research criteria and the data was analyzed by using SPSS V-16 software.

**Results:** The results showed that CSR was more observed between the ages of (26-35) (56%), and were mainly housewives (71.3%), matriculated (47.8%), and belonged to the Balochis family. As, CSR has drastically increased in Lyari Town Karachi, Pakistan (33%), and the rate of Planned Cesarean Section (CS) (63%) exaggerated more than an Emergency CS (37.4%). Scarred Uterus (32.1%) being an obstetrical factor, while Patient Preferences (30.4%) being a non-obstetrical factor were the most common reasons for Planned C-Section. Fetal Distress (33.9%) and Polyhydramnios (11.3%) were the reasons for an Emergency C-Section. Labor anxiety also influenced cesarean section. Besides, smokeless tobacco (Pan, Ghutka, Chalia, and Supari), (18% to 23%) was also identified as contributing factor among women.

**Practical Implication:** The study found a strong correlation between Corporate Social Responsibility (CSR) and demographic factors and medical practices, particularly among women aged 26-35. In Karachi, Pakistan, CSR activities increased significantly. Women prefer Planned Cesarean Sections over Emergency Cesarean Sections, with scarred uterus and patient preferences being primary reasons.

**Conclusion:** Planned CS is increasing unnecessarily in the health care setting, and this could be reduced by offering written guidelines, policy development, monthly medical audits, and advised CS when there is a medical necessity.

**Keywords:** Post-Partum Women, Planned Cesarean Section, Emergency Cesarean Section, Scarred Uterus, Fetal Distress.

## INTRODUCTION

Caesarean Section is the most common and considered safe and preferable obstetric procedure when done electively and as well as it's also a life-saving surgical procedure<sup>1</sup>. As, according to World Health Organization WHO it suggested that CSR should not exceed 10% to 15% unless any medical complication will arise<sup>1</sup>. However, CSR steadily rising in both developed and non-developed countries regardless of any medical morbidity<sup>1</sup>. Similarly, this evidence is also supported by the report of the Organization for Economic Corporation and Development Countries (OECD) that CSR has been increasing, ranging from 14.9% (Israel) to 54.9% (Turkey)<sup>2</sup>. There is no difference in the Pakistan context that the incidence of elective CSR is gradually increasing (69.7%) in public health care in Multan, Pakistan<sup>3</sup>. This study is also correlated to another study about the incidences of CSR and it observed more in Punjab (69.7%) than Sindh (17.4%), Khyber Pakhtunkhwa (KPK) (5.3%), and Baluchistan (1.70%)<sup>4</sup>.

In addition, one of the retrospective studies done from 1<sup>st</sup> January to 30<sup>th</sup> June 2018 at Jinnah Postgraduate Medical Centre (JPMC), in Karachi, Pakistan to evaluate the rates and factors of C-Section amongst pregnant women and the data of six months were retrieved and it's showed that the rate of C-Section was observed more 760(61.2%) in multigravidas than 482(38.8%) women who had delivered the first or second baby by surgically<sup>5</sup>.

Likewise, CSR was observed in the same context in Fauji Foundation Hospital (49%), and Pak Emirates Military Hospital (54%) than in JPMC (36.5%) and Holy Family Hospital, Karachi (37%)<sup>5</sup>. The reasons for escalating the numbers of C-Sections globally were; Premature Rupture Membrane (PROM), Cephalic

Pelvic Disproportion (CPD), Prolonged Fatal Distress, Multiple Pregnancies, Breech or Transverse presentation, Ante-Partum Haemorrhage (APH), Mother with HIV infection, Maternal Preference, birth weight, and limited visits antenatal<sup>5,6</sup>.

As well as, socioeconomic status such as; age, social class, education, occupation, and type of residence has been found to be strongly correlated reasons for CSR<sup>4,5,7</sup>. In addition, the authors disclosed that apart from medical factors, non-medical factors were also identified in one of the Asian countries and this qualitative study was done in both private and public sectors in Punjab, Pakistan and as cited that "perception about previous C-section, shared an experience with others women in the social circle, "family perception that vaginal delivery had not appropriate for our family, fear about vaginal delivery that it lose the beauty of the skin and its more painful than C-Section, women with high socioeconomic status and those who were not aware of the mode of delivery and this life threatening procedure its only done for the earning purposes or incentives by the gynecologist and its organization<sup>5</sup>.

Hence, these kinds of feelings changed women's perceptions of normal delivery and moved their pregnancy finish up by the surgical deliveries. According to the restating literature CSR has been increasing worldwide especially in at urban areas<sup>9</sup>. However, no studies have been conducted in densely populated underprivileged urban areas where people of different cultures are residing with low socioeconomic status.

Therefore, this study had conducted in two private tertiary care hospitals, who are providing maternity services from 1918 and 1924 to their surrounding community<sup>10,11</sup> to determine the prevalence and associated factors of CSR in two private hospitals at Lyari town Karachi, Pakistan

Received on 21-12-2023

Accepted on 26-02-2024

## METHODOLOGY

A descriptive Cross-sectional study was conducted in two private hospitals in Lyari Town Karachi, Pakistan from August to November 2022 and purposive sampling technique was used to achieve a sample size of 230. All post-partum women with age <45 were included in the study whereas women who were delivered vaginally, and those who were not willing were excluded in this research criteria. The approval was taken from the Ethical Review Committee of the Kharadar General Hospital. Written informed consent was also taken before data collection, and also maintained anonymity, and confidentiality throughout the study period. A tool of Self- Structure Questionnaire was checked by the IRB of the Kharadar General Hospital, and a panel of three experts who had maternity specialty before using it to ensure that all questions were consistently conveyed and carried the anticipated meaning of the study, and it used for data collection and data was analyzed by using SPSS V-16. Though, quantitative variables frequencies and percentages are presented in the form of tables.

## RESULTS

**Socio-Demographic Domain:** According to the Socio-demographic domain, the majority of patients were in between 26-35 years 128(55.7%), matriculated 110(47.8%), Muslims 178(77.4%) and belonged to the Balochi community 61(26.5%). In addition, data also revealed that these women were from the extended family 137(59.6%), were mainly house wives 164(71.3%), and were non-addictive 110(48%) then as compared to Chalia/Supari 54(23%), Pan/Ghutka 41(18%), Cigarette/ Berri 22(10%) and alcoholic 3(1%) were identified (Table 1).

Table 1: socio Demographic domain

Variables	Characteristics	Frequency	%age
Mother age (years)	17-25	72	31.3
	26-35	128	55.7
	36-45	30	13.0
	46-55	10	4.3
Education	Non educated	66	28.7
	Primary	11	4.8
	Secondary	6	2.6
	Matriculation	110	47.8
	Intermediate	28	12.2
	Graduation	9	3.9
Religion	Islam	178	77.4
	Hinduism	24	10.4
	Christian	27	11.7
	Others	1	0.5
Ethnicity	Memon	56	24.3
	Balochi	61	26.5
	Sindhi	44	19.1
	Punjabi	36	15.7
	Pathan	21	9.1
	Hindustani	12	5.3
Type of family	Nuclear family	93	40.4
	Extended family	137	59.6
Profession	House wife	164	71.3
	Working women	66	28.7
Level of income	Less than 20k	28	12.2
	20K-40K	115	50.0
	40K-60K	14	6.1
	60K-100K	35	15.2
	More than 100k	38	16.5
Feelings of craving or addicting	Pan/Ghutka	41	18
Chalia/Supari	Chalia/Supari	54	23
	Cigarette/Bheri	22	10
	Alcohol	3	1.0
	Mo addiction	110	48

**Clinical Domain:** In the clinical domain, most participants 99, 43%) were visited by their doctors about 11-15 times and their weights were between 50-60kg 122(53%). Furthermore, the results of our study also revealed that most of the participants had

undergone to Planned C-Section 144(62.6%) as compared to an emergency C-Section 86(37.4%), and a majority had their second pregnancies 83(36%). As regards to Planned C-Section, data also revealed that scarred Uterus was the most leading cause of caesarean section 74(32.1%) as compared to mothers' preferences 70(30.4%), malposition 24(10.4%), Intra Uterine Growth Retardation 19(8.5%), Large Fetus 25(10.8%), Multiple Fetus 15(6.5%), and macrosomia 3(1.3%). On the other hand, in emergency C-Section fetal distress 78(33.9%) and Polyhydramnios 26(11.3%) were identified the main cause, as compared to other related causes of an emergency C-Section (Table 2).

Table 2: Clinical domain

Variables	Characteristics	Frequency	%age	
Antinatal Visits	Less than 5	17	7.4	
	6-10 visits	90	39.1	
	11-15 visits	99	43.0	
	More than 15 visits	24	10.5	
Patient's current weight	Below 50kg	17	7.4	
	50-60kg	122	53.0	
	60-70kg	70	30.4	
	More than 70kg	21	9.2	
Mode of delivery	Planned CS	144	62.6	
	Emergency CS	86	37.4	
Type of pregnancy	Primi Gravida	66	28.6	
	Second Gravida	83	36.0	
	Multi gravid (1-3)	68	29.5	
	Grand Multi Gravida >4	13	5.9	
Reason of planned CS	Multiple fetus	15	6.5	
	Malposition	24	10.4	
	Large fetus	25	10.8	
	Mother's preference	70	30.4	
	Scarred uterus	74	32.1	
	Intra uterine growth	19	8.5	
	Retardation (IUGR)	3	1.3	
	Macrosomia			
Reason of emergency CS	Bad presentation	10	4.3	
	Preterm delivery	16.5	7.1	
	Pre mature Rupture of Membrane (PROM)	15	6.5	
	Polyhydramnios	26	11.3	
	Oligohydramnios	10	4.3	
	Fetal distress	78	33.9	
	Diabetes status	10	4.3	
	Prolonged obstructed labor	12.5	5.4	
	Eclampsia Failed VBAC trial	10.5	4.5	
	Cord prolapse	8	3.4	
	Placenta previa	15	6.5	
	Antepartum	11.5	5	
Haemorrhage (APH)	7	3.5		

**Socio-Psychological Domain:** Most participants 136(59.1%) performed their surgical deliveries at their own request. Labor anxiety 181(78.7%) also contributed to cesarean section, whereas reasons for caesarean section are not supported by the enforcement of their relatives 225(97.8%), financial instability 126(54.8%) and a prolonged period of hospitalization 189(82.2%) (Table 3).

Table 3: Socio Psychological domain

Variables	Characteristics	Frequency	%age
Patient request	Yes	136	59.1
	No	94	40.9
Enforced by relative	Yes	5	2.2
	No	225	97.8
Labour anxiety	Yes	181	78.7
	No	49	21.3
Financial overburdened.	Yes	104	45.2
	No	126	54.8
Extended period of hospitalization	Yes	41	17.8
	No	189	82.2

## DISCUSSION

According to another study, results showed that planned CS has increased due to scarred uterus (50%), as this result is consistent with the recent study that planned CS has increased due to the reason of previous CS (62.6%)<sup>8,13,15-19</sup>. Pertaining to the recent study, results also indicated that planned CS has increased due to patient's preferences (30.4%) as this result is also similar to the another study<sup>6,13,15,18-20</sup>. With regards to the results of the current study, that emergency CS has increased due to fetal distress (33.9%), as this result is congruent with another study that the rate of emergency CS was more observed due to fetal distress (33.5%)<sup>6,17</sup>.

Furthermore, labor anxiety also contributed to the caesarean section and most women from urban areas made a choice of delivery as CS to keep themselves away from the painful outcome and also were concerned about the safety of the baby and unfavorable procedures during labor. While this study is related with the previous study that anxiety during labor leads women towards the CS<sup>20,21</sup>.

Similarly, one of the cross-sectional studies conducted in the department of Obstetrics and Gynecology Pak Emirates Military Hospital, Rawalpindi from Jan 2018 to Jun 2018 where the authors revealed that fear of maternal in the third trimester is relatively high than the first and second trimester and it enforced by maternal to have the CS because of intense fear of childbirth.<sup>22,23</sup> Also this study conducted in Sakarya University Education and Research Hospital, Turkey where Fear of Childbirth and Post-Partum Period Scale (FCPPS) was used to identify anxiety during childbirth and results are indicated that patients who are 365(73%) unemployed 299(59.8%) and graduated, are moving towards the CS because of intense fear of childbirth during in third trimester. Similarly, one of the cross-sectional studies conducted in the Department of Obstetrics and Gynecology at Pak Emirates Military Hospital, Rawalpindi, from January 2018 to June 2018 revealed that fear of childbirth is significantly higher in the third trimester compared to the first and second trimesters. This increased fear among expectant mothers during the later stages of pregnancy often leads to a preference for cesarean section (CS) due to intense anxiety about the birthing process. The study underscores the importance of addressing maternal fears to potentially reduce the rate of elective CS<sup>18,19,20</sup>.

Additionally, a similar study conducted at Sakarya University Education and Research Hospital in Turkey utilized the Fear of Childbirth and Post-Partum Period Scale (FCPPS) to assess anxiety levels during childbirth. The findings indicated that a significant proportion of patients, 365(73%) unemployed and 299(59.8%) graduated, exhibited a preference for CS due to intense fear of childbirth in the third trimester. These results highlight the crucial role that socioeconomic and educational factors play in maternal anxiety and decision-making regarding childbirth methods<sup>21,22</sup>.

## CONCLUSION

The study concludes that C-Section Rate has drastically increased (33%) in both hospitals in Lyari Town Karachi, Pakistan. As prevalence of Planned C-Section (Scarred Uterus and Patients Preferences) seemed higher than an Emergency C-Section (fetal distress and Polyhydramnios)

**Limitations:** The limitation of this research study is that it was done only in private hospitals, as approval from public hospitals could not be obtained due to time constraints. Part from this, the research was only conducted in two private hospitals in Lyari town Karachi, and therefore, this study could not be generalized.

**Recommendations:** The result of this study suggests Written guidelines and policy development for the health sector, monthly medical audits in both public and private hospitals, commitment of Obs. /Gynae doctors to lower the C-section rate, C-Section advised when there is medical necessity, and provides antenatal

teaching to aware of the outcome of a delivery and to cope with painful delivery by reducing their anxiety in their third trimester

**Acknowledgement:** The authors would like to thanks to Mr: Zoraiz Hussain Faculty of Information Technology and Administrative Members of School of Nursing, Kharadar General Hospital Karachi who participated in this study

**Funding statement:** This research received no supporting funds from any funding agency in the public, commercial, or not-for-profit sector and it was completely done by the authors itself.

**Conflict of Interest:** The authors declared of no conflict of interest.

**Ethics approval statement:** The approval was taken from both Hospital for data collection.

**Authorship Contribution:** SP conceptualized the project, SB worked in data Collection and literature search by EF. MY performed the statistical analysis. While, Writing, revision, and writing of manuscript done by TA, SK, & SP.

## REFERENCES

1. Ajah LO, Ibekwe PC, Onu FA, Onwe OE, Ezeonu TC, Omeje I. Evaluation of clinical diagnosis of fetal distress and perinatal outcome in a low resource Nigerian setting. *Journal of clinical and diagnostic research: JCDR*. 2016 Apr;10(4):QC08.
2. Razeq NM, Khader YS, Batieha AM. The incidence, risk factors, and mortality of preterm neonates: A prospective study from Jordan (2012-2013). *Turkish journal of obstetrics and gynecology*. 2017 Mar;14(1):28.
3. Kurji Z, Premani ZS, Mithani Y. Analysis of the health care system of Pakistan: lessons learnt and way forward. *J Ayub Med Coll Abbottabad*. 2016;28(3):601.
4. Rasool MF, Akhtar S, Hussain I, Majeed A, Imran I, Saeed H, Akbar M, Chaudhry MO, Rehman AU, Ashraf W, Alqahtani F. A cross-sectional study to assess the frequency and risk factors associated with cesarean section in Southern Punjab, Pakistan. *International journal of environmental research and public health*. 2021 Aug 21;18(16):8812.
5. Mumtaz S, Bahk J, Khang YH. Rising trends and inequalities in cesarean section rates in Pakistan: Evidence from Pakistan Demographic and Health Surveys, 1990-2013. *PLoS one*. 2017 Oct 17;12(10):e0186563.
6. Majid E, Kulsoom S, Fatima S, Zuberi BF. To evaluate rising caesarean section rate and factors contributing to it by using Modified Robson's Criteria at a tertiary care hospital. *Pakistan Journal of Medical Sciences*. 2022 Sep;38(7):2021.
7. Abebe FE, Gebeyehu AW, Kidane AN, Eyassu GA. Factors leading to cesarean section delivery at Felegehiwot referral hospital, Northwest Ethiopia: a retrospective record review. *Reproductive health*. 2015 Dec;13:1-7.
8. Karlström A, Nystedt A, Johansson M, Hildingsson I. Behind the myth—few women prefer caesarean section in the absence of medical or obstetrical factors. *Midwifery*. 2011 Oct 1;27(5):620-7.
9. Latifnejad-Roudsari R, Zakerihamid M, Merghati-Khoei E, Kazemnejad A. Cultural perceptions and preferences of Iranian women regarding cesarean delivery. *Iranian journal of nursing and midwifery research*. 2014 Feb 1;19(7 Suppl 1):S28-36.
10. Mohammed NB, Kakal F, Somani M, Zafar W. Non-invasive prenatal determination of fetal RhD genotyping from maternal plasma: a preliminary study in Pakistan. *Journal of the College of Physicians and Surgeons Pakistan*. 2010;20(4):246.
11. ShahzadiNeelum, Javeria Saleem, Sumayya, Saeeda Safi, Noor-UI-Amina, Mehwish. (2022). Clinical Outcomes of Post Placental Intrauterine Contraceptive Device Insertion in Women Delivering by Cesarean Section. *Pakistan Journal of Medical & Health Sciences*, 16(09), 304. <https://doi.org/10.53350/pjmhs22169304>
12. Alwan MM, Al-Mahdawi MA. Molecular Typing of Bacterial Pathogens Isolated from Air, Non-Living Surfaces and Clinical Sources in Diyala Governorate Using MLST Technique. *Pakistan Journal of Medical & Health Sciences*. 2022;16(11):381-.
13. Naseeb S, Bibi F, Rashid S, Ali BB. Frequency and Outcome of Full Term Pregnant Patients with Previous Cesarean Section. *Pakistan Journal of Medical & Health Sciences*. 2022 Apr 5;16(02):919-.
14. Eide KT, Morken NH, Børøe K. Maternal reasons for requesting planned cesarean section in Norway: a qualitative study. *BMC pregnancy and childbirth*. 2019 Dec;19:1-0.
15. Neelum S, Saleem J, Sumayya SS, Noor-UI-Amina M. Clinical Outcomes of Post Placental Intrauterine Contraceptive Device

- Insertion in Women Delivering by Cesarean Section. *Pakistan Journal of Medical & Health Sciences*. 2022 Oct 27;16(09):304-.
16. Junge C, von Soest T, Seidler A, Eberhard-Gran M, Garthus-Niegel S. Severe recalled labor pain and elective cesarean section in a subsequent delivery: a cohort study of Norwegian parous women. *Acta Obstetrica et Gynecologica Scandinavica*. 2021 Sep;100(9):1678-87.
  17. Zeino S, Carbillon L, Pharisien I, Tigaizin A, Benchimol M, Murtada R, Boujenah J. Delivery outcomes of term pregnancy complicated by idiopathic polyhydramnios. *Journal of Gynecology Obstetrics and Human Reproduction*. 2017 Apr 1;46(4):349-54.
  18. Parveen F, Aijaz S, Aijaz W, Qureshi N. Frequency of Obstetrical Hysterectomy in Morbidly Adherent Placenta in Women with Previous Cesarean Section in Tertiary Care Hospital. *Pakistan Journal of Medical & Health Sciences*. 2023 May 12;17(03):534-.
  19. Tadevosyan M, Ghazaryan A, Harutyunyan A, Petrosyan V, Atherly A, Hekimian K. Factors contributing to rapidly increasing rates of cesarean section in Armenia: a partially mixed concurrent quantitative-qualitative equal status study. *BMC pregnancy and childbirth*. 2019 Dec;19:1-0.
  20. Khowaja B, Mughal FB, Valliani K. The Factors Influencing Cesarean-Section Rates-A Narrative Review from Pakistan. *Pakistan Journal of Medical Research*. 2021 Oct 25;60(3):143-7.
  21. Nguyen AD, Duong GT, Do DT, Nguyen DT, Tran DA, Phan TT, Nguyen TK, Nguyen HT. Primary cesarean section rate among full-term pregnant women with non-previous uterine scar in a hospital of Vietnam. *Heliyon*. 2022 Dec 1;8(12).
  22. Zengin H, Bafali I, Caka S, Tiryaki O, Cinar N. Childbirth and postpartum period fear and the related factors in pregnancy. *Journal of the College of Physicians and Surgeons Pakistan*. 2020;30(2)

---

**This article may be cited as:** Ali SPS, Yasin M, Khan SA, Bhatti SR, Fatima e, Aziz T: Determine the Prevalence and Factors Contributing to Increase the Caesarean Section Rate amongst post-partum women at Lyari Town Karachi, *Pakistan Pak J Med Health Sci*, 2024;18(3):12-15.