

# Frequency of Oral Squamous Cell Carcinoma of Tongue in patients presenting to Ayub Teaching Hospital Abbottabad

UMAIR AKRAM<sup>1</sup>, RASHIDA HILAL<sup>2</sup>, RAHEEL HASSAN<sup>3</sup>, ABID BUKHARI<sup>4</sup>

<sup>1</sup>Registrar OMFS Dentistry Section, Abbottabad International Medical Institute, Abbottabad

<sup>2,3</sup>Lecturers, Dental Section, Ayub Medical College, Abbottabad

<sup>4</sup>Dental Surgeon, DHQ Hospital Kotli, AJK

Correspondence to Dr. Rashida Hilal, E-mail: [dr.rashida.hilal@gmail.com](mailto:dr.rashida.hilal@gmail.com), Cell: 0300-4617290

## ABSTRACT

**Aim :** To determine the frequency of oral squamous cell carcinoma of tongue in patients presenting at Ayub Teaching Hospital Abbottabad.

**Study Design:** Cross- Sectional (Descriptive) Study.

**Place and duration of study:** Department of Dentistry, MTI-ATH Abbottabad from 24<sup>th</sup> December 2020 to 25<sup>th</sup> June 2021.

**Methodology:** One hundred and sixty-six patients of oral squamous cell carcinoma of oral cavity were enrolled. Patients of age 20-80 years of age, both genders, confirmed after histopathological examination (biopsies) were included. Patients not confirmed by biopsy and squamous cell carcinoma of other intra oral sites were excluded.

**Results:** There were 126(75.9%) males and 40(24.1%) females with mean age 56.61±9.50 years and mean duration of disease 3.64±1.56 years. 22(13.3%) patients had oral squamous cell carcinoma of tongue.

**Practical implication:** The benefit of the present study is highlighting the alarming increasing incidence of oral tongue squamous cell carcinoma during the last decades, specifically in women and younger patients without the traditional risk factors of alcohol or tobacco use.

**Conclusion:** Oral squamous cell carcinoma is prevalent among males and people of higher age group with only 13.3% having tongue as primary site.

**Key words:** Oral squamous cell carcinoma, Tongue, Buccal mucosa

## INTRODUCTION

Oral squamous cell carcinoma is one of the fastest growing cancer across the globe. Approximately about 300 new cases of oral-squamous-cell-carcinoma (OSCC) arise every year, making it the 6<sup>th</sup> most common type of cancer in the world. Adult population (with age greater than 50 years) are more vulnerable towards this disease. However, recent studies also reveal that frequencies among younger patients are also on a surge<sup>1</sup>.

Oral squamous cell carcinoma in oral cavity is the most common type of malignancy, with tongue being the most affected site. Usually this disease has affected males preferentially more than the females, but now cases in females are increasing. The causative factors for oral squamous cell carcinoma commonly include alcohol, tobacco, poor oral hygiene, human papillomavirus, genetic factors, exposure to sunrays and betel nut chewing<sup>2</sup>.

For the diagnosis of this disease, tissue biopsy is used. Prognostic features, like the tumor depth, is determined from resection specimens, but are seldom studied in biopsy<sup>3</sup>.

The targeted site for OSCC is gingivobuccal mucosa (38.21%), followed by buccal mucosa (30.90%), tongue (12.29%), alveolar mucosa (8.64%), lip (3.99%), floor of mouth and hard palate. In the Western world tongue and floor of mouth carcinomas are more common due consumption of alcohol and smoking<sup>4,5</sup>. Frequency of OSCC differs in different regions of world, depending upon the mode of intake and placement trend of tobacco at different intra oral sites e.g. in Pakistan mostly snuff is placed into gingivobuccal sulcus and buccal mucosa. Thereby, making buccal mucosa the most common site followed by lips, tongue and gingiva<sup>7</sup>. Similarly, in Sri Lanka the buccal mucosa is the common site<sup>6</sup>.

The recognition of squamous cell carcinoma of tongue is increasing around the world. As no such study has been done before in our general population, therefore I have planned to determine the frequency of squamous cell carcinoma of tongue in patients presenting to Ayub Teaching Hospital, Abbottabad. The results of my study will estimate the real burden of this morbidity in our general population.

Received on 07-12-2023

Accepted on 25-02-2024

## MATERIALS AND METHODS

This cross-sectional study was conducted in the Department of Dentistry, MTI-ATH Abbottabad from 24<sup>th</sup> December 2020 to 24<sup>th</sup> June 2021 and after IRBB permission 166 patients of oral squamous cell carcinoma were enrolled. Patients of age 20-80 years, both genders, confirmed after histopathological examination (biopsy) and new cases as well as with recurrence were included. Patients not confirmed by biopsy, squamous cell carcinoma of other intra oral sites and biopsy report from untrusted sources were excluded. Biopsy was performed under supervision of consultant maxillofacial surgeon. Specimens were sent to pathology laboratory Ayub Medical College Abbottabad and examined by consultant pathologist of Ayub Medical College and results were recorded. Data was entered and analyzed with SPSS-22.

## RESULTS

There were 43(25.9%) patients ≤50 years of age while 123(74.1%) patients >50 years of age. 126(75.9%) patients were males and 40(24.1%) patients were females (Table 1). The mean age was 56.61±9.50 years and mean duration of disease was 3.64±1.569 years (Table 2). 22(13.3%) patients had oral squamous cell carcinoma of the tongue (Table 3). When the oral squamous cell carcinoma was compared with duration of disease, statistically the difference was not significant (P>0.05) [Table 4].

Table 1: Demographic information of the patients (n=166)

Variable	No.	%
<b>Age (years)</b>		
≤ 50	43	25.9
> 50	123	74.1
<b>Gender</b>		
Male	126	75.9
Female	40	24.1

Table 2: Descriptive statistics of the patients (n=166)

Variable	Mean±SD
Age (years)	56.61±9.50
Duration of disease (years)	3.64±1.56

Table 3: Frequency of Oral squamous cell carcinoma of tongue (n=166)

Variables	No.	%
Yes	22	13.3
No	144	86.7

Table 4: Squamous cell carcinoma was compared with duration of disease

Oral squamous cell carcinoma	Duration of Disease		P value
	≤ 3 years	>3 years	
Yes	11 (6.6%)	12 (7.3%)	0.626
No	80 (48.2%)	63 (37.9%)	

## DISCUSSION

Oral squamous cell carcinoma is one of the most prevalent malignancies that affect people of different age groups. Staging of cancer is crucial for planning and also for treatment planning. Late presentation, lack of awareness and diagnostic markers are the key main hurdles for treating and combating this disease. Present study was designed to determine the frequency of OSCC of tongue in patients presenting to Ayub Teaching Hospital, Abbottabad.

In this study, mean age was 56.61±9.50 years and duration of disease was 3.64±1.56 years (Table 1). 43(25.9%) patients were recorded in ≤50 years of age group while 123(74.1%) patients were recorded in >50 years of age group. 126(75.9%) male patients and 40 (24.1%) female patients (Table 2). Various studies conducted in different regions of the world presented with similar presentation<sup>7-9</sup>. A study conducted in Japan also highlights that people of >50 years of age are more vulnerable towards this disease and male were more commonly affected.<sup>8</sup> Another study conducted in China also showed the similar pattern of male to female ratio and age group<sup>9</sup>.

The frequency of oral squamous cell carcinoma differs in different regions of world, depending upon the mode of intake and placement trend of tobacco at different intra oral sites.<sup>10</sup> In the Western world tongue and floor of mouth carcinomas are more common due consumption of alcohol and smoking.<sup>5,14,15</sup> In our study only 22(13.3%) patients had oral squamous cell carcinoma of the tongue (Table 3) as in Pakistan snuff is most commonly placed in the gingivobuccal sulcus and the buccal mucosa. Similarly, a study conducted in Sri Lanka showed buccal mucosa being the most favoured site for OSCC<sup>6</sup>. This study also demonstrated that highest proportion of patients 14(11.4%) were recorded in >50 years' age group in our local population and risk of OSCC also get increased with the passage of age and time.

## CONCLUSION

Squamous cell carcinoma is a very common malignancy of the oral cavity which is more prevalent in males and people of higher age group. Compared to the Western world, tongue was not as frequently involved in our region according to this study. However, multicentre studies are required to estimate the frequency of involvement of different intraoral sites and the disease burden of OSCC.

**Authorship and contribution declaration:** Each author of this article fulfilled following Criteria of Authorship:

1. Conception and design of or acquisition of data or analysis and interpretation of data.

2. Drafting the manuscript or revising it critically for important intellectual content.
3. Final approval of the version for publication.  
All authors agree to be responsible for all aspects of their research work.

**Funding:** None

**Conflict of interest:** The authors declare no conflict of interest in this study.

## REFERENCES

1. Migueláñez-Medrán BC, Pozo-Kreilinger JJ, Cebrián-Carretero JL, Jinez-García MA, Lopez-Sánchez AF. Oral squamous cell carcinoma of tongue: Histological risk assessment: a pilot study. *Med Oral Patol Oral Cir Bucal* 2019;24(5):24(5):e603-9.
2. Zafereo ME, Xu L, Dahlstrom KR, Viamonte CA, El-Naggar AK, Wei Q, et al. Squamous cell carcinoma of the oral cavity often over-expresses p16 but is rarely driven by human papillomavirus. *Oral Oncol* 2016; 56:47-53.
3. Dhanda J, Uppal N, Chowlia H, Opie N, Al-Qamachi L, Shelat D, et al. Features and prognostic utility of biopsy in oral squamous cell carcinoma. *Head Neck* 2016; 38: E1857-62.
4. Smitha T, Mohan CV, Hemavathy S. Clinicopathological features of oral squamous cell carcinoma: A hospital-based retrospective study. *J NIR Univ Health Sci* 2017;6:29-34.
5. Singh MP, Kumar V, Agarwal A, Kumar R, Bhatt ML, Misra S. Clinic-epidemiological study of oral squamous cell carcinoma: a tertiary care centre study in North India. *J Oral Biol Craniofac Res* 2016;6(1):31-4.
6. Jayasooriya PR, Pitakotuwage TN, Ranjit B, Mendis RN, Lombardi T. Descriptive study of 896 Oral squamous cell carcinomas from the only University based Oral Pathology Diagnostic Service in Sri Lanka. *BMC Oral Health* 2016; 16: 10.
7. Imtiaz AS, Hurtamina K, Riaz S, Mahum H, Muhammad RD, et al. Oral cancer frequency at different sub sites presenting at a tertiary care hospital in Karachi Pakistan. *Glob J Oto* 2017; 6(3): 555687.
8. Vigneswaran N, Williams. Epidemiologic trends in head and neck cancer and aids in diagnosis. *Oral Maxillofac Surg Clin North Am* 2014; 26:123-41.
9. Zhong LP, Zhou XJ, Wei KJ, Yang X, Ma CY, Zhang CP, et al. Application of serum tumor markers and support vector machine in the diagnosis of oral squamous cell carcinoma. *Shanghai Kou Qiang Yi Xue* 2008;17(5):457-60.
10. Vanderveken OM, Szturz P, Specenier P, Merlano MC, Benasso M, Van Gestel D, et al. Gemcitabine-based chemoradiation in the treatment of locally advanced head and neck cancer: systematic review of literature and meta-analysis. *Oncologist* 2016; 21(1): 59-71.
11. Chinn SB, Myers JN. Oral cavity carcinoma: current management, controversies, and future directions. *J Clin Oncol* 2015; 33(29):3269-76.
12. Nwizu T, Adelstein D. Pharmacotherapy of head and neck cancer. *Expert Opin Pharmacother* 2015; 16(16):2409-22.
13. Brown TJ, Gupta A. Management of Cancer Therapy-Associated Oral Mucositis. *JCO Oncol Pract* 2020; 16 (3):103-9.
14. Allison RR, Ambrad AA, Arshoun Y, Carmel RJ, Ciuba DF, Feldman E, et al. Multi-institutional, randomized, double-blind, placebo-controlled trial to assess the efficacy of a mucoadhesive hydrogel (MuGard) in mitigating oral mucositis symptoms in patients being treated with chemoradiation therapy for cancers of the head and neck. *Cancer* 2014; 120 (9):1433-40.
15. Hong CHL, Gueiros LA, Fulton JS, Cheng KKF, Kandwal A, Galiti D, et al. Systematic review of basic oral care for the management of oral mucositis in cancer patients and clinical practice guidelines. *Support Care Cancer* 2019; 27(10): 3949-67.

**This article may be cited as:** akram U, Hilal R, Hassan R, Bukhtari A: Frequency of Oral Squamous Cell Carcinoma of Tongue in patients presenting to Ayub Teaching Hospital Abbottabad. *Pak J Med Health Sci*, 2024;18(3):27-28