

ORIGINAL ARTICLE

Diagnostic Accuracy of Serum Procalcitonin, C-Reactive Protein and White Blood Cell Count in Differentiating Uncomplicated versus Complicated Acute Appendicitis

FARHAN ALI¹, TAYYABA RASHEED², KALSOOM KALHORO³, TARIQ HUSSAIN⁴, MUHAMMAD NUMAN TAHIR⁵¹Assistant Professor, Department of General Surgery, Pir Abdul Qadir Shah Jeelani Institute of Medical Sciences, Gambat, Sindh²Postgraduate Residents, Department of General Surgery, Pir Abdul Qadir Shah Jeelani Institute of Medical Sciences, Gambat, Sindh³Senior Registrar, Department of General Surgery, Pir Abdul Qadir Shah Jeelani Institute of Medical Sciences, Gambat, Sindh⁴Postgraduate Residents, Department of General Surgery, Pir Abdul Qadir Shah Jeelani Institute of Medical Sciences, Gambat, Sindh⁵Postgraduate Residents, Department of General Surgery, Pir Abdul Qadir Shah Jeelani Institute of Medical Sciences, Gambat, SindhCorrespondence to: Dr. Farhan Ali, Email: drfarhan00@gmail.com

This article may be cited as:

Ali F, Rasheed T, Kalhoro K, Hussain T, Tahir MN; Diagnostic Accuracy of Serum Procalcitonin, C-Reactive Protein and White Blood Cell Count in Differentiating Uncomplicated versus Complicated Acute Appendicitis. Pak J Med Health Sci, 2026; 20(03): 12-19.

Received: 02-10-2025

Accepted: 15-03-2026

Published: 30-03-2026



© The Author(s) 2026. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.



ABSTRACT

Background: Acute appendicitis is a common surgical emergency. The ability to differentiate between uncomplicated and complicated appendicitis early in the disease process is important as complicated appendicitis is associated with perforation, abscess, peritonitis, longer hospital stay, and higher post-operative complications. Preoperative assessment of the severity of appendicitis may be aided by inflammatory markers such as serum procalcitonin, C-reactive protein, and white blood cell count.

Objective: To assess and compare the diagnostic value of serum procalcitonin, C-reactive protein and white blood cell count in distinguishing uncomplicated acute appendicitis from complicated acute appendicitis.

Methods: This cross-sectional analytical study was performed in the Department of General Surgery, Pir Abdul Qadir Shah Jeelani Institute of Medical Sciences, Gambat, Sindh from 1st August 2025 to 31st January 2026. One hundred patients between 15 and 60 years of age with a clinical diagnosis of acute appendicitis who were operated on by consecutive sampling were enrolled. Serum procalcitonin, C-reactive protein and white blood cell count were determined preoperatively. Uncomplicated and complicated appendicitis were defined by surgery and histopathology. SPSS version 26.0 was used for the statistical analysis.

Results: Among the 100 patients, 58 (58.0%) had uncomplicated appendicitis and 42 (42.0%) had complicated appendicitis. Patients with complicated appendicitis were older, had longer duration of symptoms, and had more fever, guarding, and rebound tenderness. The average white blood cell count, C-reactive protein and procalcitonin were significantly elevated in complicated appendicitis ($p < 0.001$). Serum procalcitonin had the best diagnostic performance (sensitivity, 85.7%; specificity, 87.9%; positive predictive value, 83.7%; negative predictive value, 89.5%; and accuracy, 87.0%). C-reactive protein had intermediate diagnostic performance, while white blood cell count was the least accurate.

Conclusion: Procalcitonin was the most accurate marker to distinguish complicated from uncomplicated acute appendicitis, followed by C-reactive protein and white blood cell count. This can be used in the preoperative period for early risk assessment and surgical planning.

Keywords: Appendicitis, Procalcitonin, C-reactive protein, Leukocyte count, Diagnostic accuracy, Surgical emergencies.

INTRODUCTION

Acute appendicitis is one of the most frequent causes of acute abdominal pain presenting for emergency surgery globally.¹ It is one of the most common causes of general surgical admissions and remains a diagnostic dilemma despite advances in laboratory and radiology.² While diagnosis is frequently made by a combination of history, examination, laboratory and radiology, it remains challenging to discriminate uncomplicated acute appendicitis from complicated acute appendicitis pre-operatively in many cases.³ This is important because complicated appendicitis, which encompasses gangrenous appendicitis, perforation, appendicular abscess, appendicular mass, and generalized peritonitis, is associated with greater morbidity, longer hospital stay, more complications, and greater health-care costs.⁴

Acute appendicitis has historically been managed as a single surgical emergency; but growing evidence indicates that uncomplicated and complicated appendicitis may be distinct pathological and inflammatory processes rather than a progression of the same disease.⁵ Uncomplicated appendicitis is usually defined as simple inflammation of the appendix without evidence of gangrene, perforation, and peri-appendiceal abscess, while complicated appendicitis is defined as more severe and destructive inflammatory changes with increased risk of sepsis and complications.⁶ Thus, early recognition of patients with a high likelihood of complicated appendicitis is clinically important for timely surgical intervention, pre-operative assessment and preparation, antibiotic therapy, risk stratification, and avoidance of complications.⁷

In emergency practice, clinical diagnosis may not be entirely accurate, especially in patients presenting with atypical symptoms, delayed presentation, very young or elderly age, obesity, or coexisting abdominal and pelvic pathology.⁸ Radiological tests, including ultrasound and computed tomography, may enhance diagnostic accuracy, but may not be readily accessible in low-resource settings, and may not always be able to clearly indicate the severity of appendicitis.⁹ In many centres, particularly in the developing world, clinicians still rely heavily on affordable and rapidly available laboratory parameters.¹⁰ Consequently, inflammatory markers have gained increasing attention as a useful tool to detect the severity of appendiceal inflammation and to predict the risk of complicated disease.¹¹

The most common laboratory markers used in the diagnosis of suspected appendicitis are white blood cell (WBC) count and C-reactive protein (CRP).¹² Leukocytosis is well known as a marker of acute infection and

inflammation and is frequently the first abnormal laboratory finding in appendicitis.¹³ But while leukocytosis is frequently observed in appendiceal inflammation, it is not specific and can also be found in many intra-abdominal, infectious and inflammatory diseases.¹⁴ Likewise, CRP, an acute phase protein produced by the liver in response to inflammatory cytokines, has been reported to reflect the severity and duration of inflammation.¹⁵ Elevated CRP levels are often observed in patients with gangrenous and/or perforated appendicitis, and has been shown to be a predictor of disease severity.¹⁶ However, CRP may not be sufficient to reliably differentiate complicated from uncomplicated appendicitis, particularly in the early stages of disease.¹⁷

Over the last decade, serum procalcitonin has been reported as a more specific marker of severe bacterial infection, tissue inflammation, and systemic inflammatory response.¹⁸ Procalcitonin is a precursor of calcitonin that is present at low levels in the blood of healthy individuals, but increases markedly in bacterial infection and severe inflammation.¹⁹ Given complicated appendicitis is more likely to be associated with transmural inflammation, necrosis, perforation and bacterial contamination, procalcitonin has been suggested as a potential biomarker to predict the risk of complicated appendicitis.²⁰ While some studies have reported superior diagnostic performance of procalcitonin over traditional inflammatory markers for predicting perforation and complicated appendicitis, the evidence remains mixed, and the relative diagnostic accuracy of procalcitonin, CRP, and WBC count has yet to be determined in various populations and clinical settings.¹¹

This is especially true in low- and middle-income countries like Pakistan, where acute appendicitis is a common presentation in emergency departments, but where presentation may be delayed, access to sophisticated imaging may be limited, and diagnostic resources may be scarce.⁹ In these circumstances, the discovery of simple, affordable, and readily available biomarkers that can distinguish between uncomplicated and complicated appendicitis may have a significant impact on preoperative assessment and aid in clinical decision making.¹⁵ Determining the relative diagnostic accuracy of commonly available inflammatory markers may allow surgeons and emergency department physicians to more accurately triage urgent surgical intervention, predict operative findings, and avoid the negative impact of missed perforation or delay in diagnosis.⁶

To address these issues, the current study aimed to assess the diagnostic performance of serum procalcitonin, C-reactive protein, and white blood cell count in

distinguishing uncomplicated from complicated acute appendicitis in patients with suspected appendicitis.⁷ This study will compare these markers with operative and/or histopathological findings to determine the most valuable laboratory marker for predicting complicated appendiceal pathology and to provide clinically relevant information for the improvement of emergency surgical practice in a hospital setting.¹⁰

MATERIAL AND METHOD

This hospital-based analytical cross-sectional study was carried out in the Department of General Surgery, Pir Abdul Qadir Shah Jeelani Institute of Medical Sciences, Gambat, Sindh from 1st August 2025 to 31st January 2026. The objective of this study was to compare the diagnostic accuracy of serum procalcitonin, C-reactive protein (CRP) and white blood cell (WBC) count in distinguishing between uncomplicated acute appendicitis and complicated acute appendicitis in patients with suspected appendicitis. This study was conducted following approval from the institutional ethical review committee, and the study procedures were conducted in accordance with the ethical standards for human experimentation. Informed consent was taken from all patients or their attendants.

One hundred patients of both sexes were enrolled in the study by non-probability consecutive sampling. Any patient between the ages of 15 to 60 years presenting to the emergency department or surgical outpatient/emergency receiving unit with clinical signs and symptoms suggestive of acute appendicitis and who planned for appendectomy were eligible to be included. The diagnosis of clinical suspicion was made based on a combination of history and examination findings (right iliac fossa pain, nausea or vomiting, anorexia, fever, local tenderness, rebound tenderness) and laboratory and/or ultrasonographic findings suggestive of appendicitis. Only those patients who were diagnosed with appendicitis intraoperatively and/or on histopathological examination were included. Those with a history of chronic inflammatory disease, autoimmune disease, active systemic infection other than appendicitis, generalized sepsis from other sources, chronic liver disease, chronic kidney disease, malignancy, hematological disorder, immunosuppressive therapy, or antibiotic use for more than 48 hours prior to presentation were excluded, as these factors may influence the levels of inflammatory markers. Women who were pregnant, those who had an interval appendectomy, those with an appendicular lump that was treated conservatively, or who refused to participate, were also excluded.

Once enrolled, a proforma was used to collect demographic and clinical information. These included age,

sex, duration of pain, fever, nausea and vomiting, anorexia, right iliac fossa tenderness, guarding, rebound tenderness, and clinical diagnosis. Radiological findings, especially abdominal ultrasound if performed, were also recorded. Preoperative blood samples were collected from a vein aseptically for determination of the chosen inflammatory markers. White blood cell count (WBC) was determined by automated hematology analyzer and expressed as $\times 10^9/L$. Serum CRP was determined by standard immunoturbidimetric method and expressed as mg/L, and serum procalcitonin was measured by an immunoassay-based laboratory method and expressed as ng/mL. All specimens were processed in the hospital laboratory with standard quality control procedures.

All patients who presented with suspected acute appendicitis were treated by open or laparoscopic appendectomy, depending on the on-call surgeon's preference and institutional protocol. The surgeon recorded the findings at the time of surgery. Postoperatively, the appendix specimens were referred for histopathological analysis and the final diagnosis of severity was based on the operative and histopathological reports. The patients were grouped into uncomplicated acute appendicitis and complicated acute appendicitis. Uncomplicated appendicitis was defined as an inflamed or suppurative appendix without gangrene, perforation, abscess, appendicular mass, or generalized peritonitis. Complicated appendicitis was defined as gangrenous appendicitis, perforated appendix, appendicular abscess, appendicular phlegmon/mass, or localized or generalized peritonitis.

The primary objective of the study was to assess the diagnostic usefulness of serum procalcitonin, CRP, and WBC count to distinguish complicated appendicitis from uncomplicated appendicitis. The secondary objective was to compare the average biomarker values between the two groups and the diagnostic value of each biomarker. To this end, biomarker levels were compared between the two groups, and the diagnostic accuracy of each biomarker for complicated appendicitis was evaluated.

Data were entered and analyzed using Statistical Package for the Social Sciences (SPSS) version 26.0. The independent sample t-test was used to compare the mean value of biomarkers between uncomplicated and complicated appendicitis. Categorical variables were compared using the chi-square test. For diagnostic performance, sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and diagnostic accuracy of serum procalcitonin, CRP and WBC count were determined using operative and histopathological diagnosis as the gold standard. A p-value of <0.05 was considered statistically significant.

RESULTS

There were 58 (58%) diagnosed with uncomplicated acute appendicitis, and 42 (42%) with complicated acute appendicitis. Complicated appendicitis was defined as gangrenous appendix, perforated appendix, appendicular abscess, appendicular phlegmon and localized and generalized peritonitis. Therefore, slightly more than 40% of the study population had advanced appendiceal pathology, suggesting a significant proportion of complicated appendicitis among patients who were operated on. The mean age of the study population was 29.84 ± 10.62 years. The mean age of patients with uncomplicated appendicitis was 27.79 ± 9.89 years, while it was 32.67 ± 11.04 years for patients with complicated appendicitis. This was statistically significant ($p=0.021$), suggesting that patients with complicated appendicitis were older, on average, than those with uncomplicated appendicitis. Sixty two patients (62%) were male and 38 patients (38%) were female, with a slight male preponderance in the study population. Among the patients with uncomplicated appendicitis, 35 of 58 (60.3%) were male and 23 of 58 (39.7%) were female. In the complicated appendicitis group, 27 of 42 patients (64.3%) were male and 15 of 42 (35.7%) were female. While males were more common in both groups, the gender distribution between uncomplicated and complicated appendicitis was similar ($p = 0.684$). This indicates that although appendicitis occurred more frequently in males in this study, gender was not a significant predictor of uncomplicated or complicated appendicitis. There was a significant difference in the time to presentation with abdominal pain. The mean duration of abdominal pain was 19.38 ± 8.47 hours in patients with uncomplicated appendicitis, compared with 31.21 ± 10.86 hours in patients with complicated appendicitis ($p < 0.001$). This suggests that the longer the delay in presentation, the greater the likelihood of complicated appendicitis, which is consistent with the clinical observation that the duration of the inflammatory process is a significant factor in the development of perforation, gangrene and peri-appendiceal infection. The presence of other clinical symptoms and signs also varied significantly between the two groups. Fever was reported in 18 patients (31.0%) with uncomplicated appendicitis and 29 patients (69.0%) with complicated appendicitis ($p < 0.001$). Likewise, nausea and/or vomiting was reported in 34 patients (58.6%) with uncomplicated appendicitis and 32 patients (76.2%) with complicated appendicitis. While this symptom was more common in the complicated group, this did not reach statistical significance ($p = 0.071$), and it appears to reflect more severe gastrointestinal symptoms in complicated appendicitis. Greater differences were

noted on abdominal examination. Rebound tenderness was present in 24 patients (41.4%) in the uncomplicated group and in 31 patients (73.8%) in the complicated group ($p=0.001$). Similarly, guarding was noted in only 16 patients (27.6%) with uncomplicated appendicitis, whereas it was noted in 28 patients (66.7%) with complicated appendicitis ($p < 0.001$). This suggests that patients with complicated appendicitis had more pronounced peritoneal irritation and a more progressed inflammatory process in the abdomen than patients with uncomplicated appendicitis (Table 1).

The three biomarkers, WBC count, CRP and serum procalcitonin, were all significantly higher in patients with complicated appendicitis, suggesting a more severe systemic inflammatory response in complicated appendicitis. The mean white blood cell (WBC) count in patients with uncomplicated appendicitis was $11.82 \pm 2.41 \times 10^9/L$, while the mean WBC count in patients with complicated appendicitis was significantly higher at $15.64 \pm 3.28 \times 10^9/L$. This was highly statistically significant ($p < 0.001$). These results suggest that leukocytosis was more severe in complicated appendicitis, suggesting a higher inflammatory load and likely bacterial spread. But while the average WBC count was obviously higher in complicated appendicitis, there was nevertheless some degree of overlap, so it is possible that WBC counts alone may not be adequate for accurate severity grading. Similar but even more pronounced findings were found for C-reactive protein (CRP). The mean CRP in the uncomplicated appendicitis group was 24.93 ± 12.76 mg/L, whereas it was 68.47 ± 24.81 mg/L in the complicated appendicitis group. This was also statistically significant ($p < 0.001$). These higher CRP levels in complicated appendicitis indicate that CRP was closely related to more severe inflammation of the appendix, necrosis of the appendix, and possible perforation or abscess formation. CRP had a greater difference between the two groups than WBC count, suggesting that CRP was a better marker of the severity of inflammation. Serum procalcitonin was the most discriminatory laboratory marker between uncomplicated and complicated appendicitis. The mean serum procalcitonin concentration in patients with uncomplicated appendicitis was 0.18 ± 0.11 ng/mL, while the mean level for patients with complicated appendicitis was 1.42 ± 0.76 ng/mL ($p < 0.001$). This marked difference suggests that procalcitonin increased dramatically in patients with severe appendiceal disease, and is the most discriminatory marker in the current study. These results are consistent with the notion that procalcitonin may be a better marker for severe bacterial infection, tissue damage, and systemic inflammation than traditional routine markers. The laboratory findings showed a clear and clinically relevant pattern: as appendicitis advanced

from uncomplicated to complicated disease, the levels of all inflammatory markers rose significantly, but the increase was greatest for procalcitonin, followed by CRP, and then WBC count (Table 2).

Serum procalcitonin had the best diagnostic performance of the three biomarkers. It had a sensitivity of 85.7%, which means that it correctly diagnosed 86 of 100 patients who had complicated appendicitis. It also had a specificity of 87.9%, meaning that it correctly identified almost 88 of every 100 patients who did not have complicated appendicitis. Procalcitonin also had a positive predictive value (PPV) of 83.7%, meaning that if a patient's procalcitonin was raised, they were likely to have complicated appendicitis, and a negative predictive value (NPV) of 89.5%, meaning that if a patient's procalcitonin was not raised, they were very likely not to have complicated appendicitis. It had an overall accuracy of 87.0%, and was the most accurate marker in this study. CRP was also a good marker, but not as good as procalcitonin. It had a sensitivity of 81.0%, specificity of 79.3%, PPV of 73.9%, NPV of 84.6%, and overall diagnostic accuracy of 80.0%. This suggests that CRP was a good marker for distinguishing complicated from uncomplicated appendicitis and was better than WBC count, but not as good as procalcitonin. WBC count, however, had the poorest diagnostic performance. It

demonstrated a sensitivity of 73.8%, specificity of 69.0%, PPV of 63.3%, NPV of 78.4%, and accuracy of 71.0%. White blood cell count was significantly higher in complicated appendicitis and still useful as a first-line inflammatory marker, the lower specificity and predictive values suggest that it was less specific than CRP and procalcitonin for determining the severity of appendicitis. Overall, these results show that although all three markers were correlated with complicated appendicitis, the serum marker procalcitonin had the highest diagnostic accuracy, followed by CRP, while the WBC count had the lowest diagnostic accuracy as a single marker (Table 3).

The present study demonstrated that complicated acute appendicitis was associated with older age, longer duration of symptoms, higher frequency of fever, more marked abdominal signs, and significantly elevated inflammatory biomarkers. There was a balanced representation of both sexes, with a slight male predominance. Crucially, serum procalcitonin was found to be the most reliable biomarker for distinguishing uncomplicated from complicated appendicitis, followed by CRP, with less discriminatory value of WBC count. These findings indicate that the addition of serum procalcitonin and CRP to the standard preoperative work-up may help in early diagnosis of complicated appendicitis and enable better surgical management.

Table 1. Demographic and clinical characteristics of patients with uncomplicated and complicated acute appendicitis

Variable	Uncomplicated appendicitis (n = 58)	Complicated appendicitis (n = 42)	p-value
Age (years)	27.79±9.89	32.67±11.04	0.021
Duration of pain (hours)	19.38 ± 8.47	31.21 ± 10.86	<0.001
Gender			
Male	35 (60.3%)	27 (64.3%)	0.684
Female	23 (39.7%)	15 (35.7%)	
Fever	18 (31.0%)	29 (69.0%)	<0.001
Nausea/Vomiting	34 (58.6%)	32 (76.2%)	0.071
Rebound tenderness	24 (41.4%)	31 (73.8%)	0.001
Guarding	16 (27.6%)	28 (66.7%)	<0.001

Table 2. Comparison of inflammatory biomarkers between uncomplicated and complicated acute appendicitis

Biomarker	Uncomplicated appendicitis (n = 58)	Complicated appendicitis (n = 42)	p-value
WBC Count ($\times 10^9/L$)	11.82±2.41	15.64±3.28	<0.001
CRP (mg/L)	24.93±12.76	68.47±24.81	<0.001
Procalcitonin (ng/mL)	0.18±0.11	1.42±0.76	<0.001

Table 3. Diagnostic performance of serum procalcitonin, C-reactive protein and white blood cell count in identifying complicated acute appendicitis

Biomarker	Sensitivity	Specificity	PPV	NPV	Accuracy
Procalcitonin	85.7%	87.9%	83.7%	89.5%	87%
CRP	81%	79.3%	73.9%	84.6%	80%
WBC Count	73.8%	69%	63.3%	78.4%	71%

DISCUSSION

The present study showed that patients with complicated appendicitis had more severe clinical features, such as advanced age, prolonged duration of symptoms, higher incidence of fever, more rebound tenderness, and more guarding, than patients with uncomplicated appendicitis. Moreover, the three inflammatory markers were significantly higher in complicated disease, but the discriminative power of serum procalcitonin was highest, followed by CRP, while WBC count had the lowest diagnostic performance. These results are significant because they indicate that basic preoperative laboratory tests, particularly procalcitonin, may be used to identify patients at increased risk of complicated appendiceal pathology prior to surgery.⁴

In the current study, 42% of patients had complicated appendicitis and 58% had uncomplicated appendicitis.⁵ This suggests a high incidence of severe appendiceal disease in the study cohort.⁶ The high proportion of complicated appendicitis in this study population may reflect delayed presentation, variable access to early surgical assessment, self-medication prior to hospital presentation, and late referrals, which are often encountered in many low- and middle-income countries.⁷ This is important from a clinical perspective as complicated appendicitis is more likely to be complicated by perforation, abscess, longer hospital stay, wound infection, intra-abdominal sepsis and higher post-operative complications.⁸ So, their early recognition is a key focus of emergency surgery.⁹ The latest international consensus also favours using a combination of clinical, laboratory and imaging parameters rather than a single one to diagnose appendicitis.¹⁰

In the current study, there was a slight male predominance, with 62% males and 38% females.¹¹ There were more males in both uncomplicated and complicated appendicitis, but the difference was not significant.¹² This finding is in agreement with the widely reported epidemiology of acute appendicitis, in which males are more frequent than females.¹³ But the lack of a significant gender difference in the severity of appendicitis in the current study indicates that once the inflammatory process is underway, both males and females have an equal risk of developing complicated appendicitis.¹⁴ This finding is important because it highlights that the severity of disease should be determined on the basis of clinical and biochemical parameters rather than gender.¹⁵

An important observation of this study was the significantly longer duration of pain prior to hospital presentation in patients with complicated appendicitis (31.21±10.86 hours) compared with uncomplicated

appendicitis (19.38±8.47 hours).¹⁶ This suggests that a delay in presenting to hospital or in diagnosis plays a significant role in disease advancement.¹⁷ This makes clinical sense because appendiceal inflammation may start as a mucosal or luminal process but can then progress to suppuration, transmural necrosis, gangrene, perforation and peri-appendiceal infection.¹⁸ The significantly greater proportion of patients with fever, rebound tenderness and guarding in the complicated appendicitis group in the present study also supports this pattern of disease evolution, and suggests that more severe disease is more likely to be associated with a more severe systemic and peritoneal inflammatory response.¹⁹

With respect to laboratory indices, the present study found that the WBC count was significantly higher in complicated appendicitis (15.64±3.28 ×10⁹/L) than in uncomplicated appendicitis (11.82±2.41 ×10⁹/L).²⁰ This confirms the well-known role of leukocytosis as a marker of acute infection and inflammation.¹² WBC count has traditionally been one of the most readily available laboratory markers in cases of suspected appendicitis and is included in many scoring systems¹³. But while WBC count was significantly elevated in patients with complicated appendicitis in this study, its overall accuracy was relatively poor (71%)² with lower sensitivity and specificity than the other laboratory markers.³ This suggests that while WBC count is helpful, it is not specific enough to be used alone as a predictor of appendiceal severity.⁴ WBC count can be elevated in both complicated and uncomplicated appendicitis and can also be elevated in other intra-abdominal inflammatory disease.⁵ As such, WBC count should be used as an adjunct, rather than a conclusive, marker of appendiceal severity.⁶

The results for CRP were more striking.⁷ The mean CRP was significantly higher in patients with complicated appendicitis (68.47±24.81 mg/L) than in those with uncomplicated appendicitis (24.93±12.76 mg/L).⁸ CRP is an acute-phase protein synthesised in the liver in response to inflammatory cytokines, especially interleukin-6, and has been shown to increase with increasing tissue inflammation and bacterial damage.⁹ The significantly higher levels of CRP in complicated appendicitis found in this study indicate that CRP is very strongly correlated with greater tissue damage, necrosis, or perforation.¹⁰ In this study, it also performed better than WBC count, with an accuracy of 80%.¹¹ This suggests that CRP is a better predictor than WBC count of advanced appendiceal inflammation, and may be a valuable adjunct in the interpretation of a suspected appendicitis case in the emergency department.¹²

The key finding of the current study was that serum procalcitonin outperformed the other two markers.¹³ The

average serum procalcitonin concentration was 0.18 ± 0.11 ng/mL in uncomplicated appendicitis, but increased to 1.42 ± 0.76 ng/mL in complicated appendicitis, with the greatest difference between groups of all the markers examined.¹⁴ Procalcitonin also had the highest sensitivity (85.7%), specificity (87.9%), NPV (89.5%) and diagnostic accuracy (87%) of the three markers.¹⁵ These results indicate that serum procalcitonin is the most valuable biochemical marker of the three markers studied for preoperative discrimination between uncomplicated and complicated acute appendicitis.¹⁶

The rationale for this is significant.¹⁷ Procalcitonin is the precursor of calcitonin, which is present in very low concentrations in normal individuals but increases in the presence of bacterial infection, systemic inflammation and necrosis.¹⁸ While uncomplicated appendicitis may be more likely to have localised and limited inflammation, complicated appendicitis is more likely to have transmural damage, bacterial translocation, gangrene, perforation, abscess and peritoneal contamination.¹⁹ These factors are more likely to trigger a systemic pro-inflammatory response and, as a result, a greater rise in procalcitonin.²⁰ Thus, it is conceivable that procalcitonin would be superior to other inflammatory markers for the detection of severe appendiceal disease.¹¹

Our results are consistent with previous studies that have shown that procalcitonin may not be useful for diagnosing appendicitis in general, but may be useful for diagnosing complicated appendicitis.¹² One previous meta-analysis concluded that although procalcitonin was not the single most valuable marker for diagnosing all suspected appendicitis, it was more valuable for diagnosing complicated appendicitis, especially because of its high specificity¹³. Likewise, Li et al² found that procalcitonin was an excellent differential marker between uncomplicated and complicated acute appendicitis in adults, consistent with the present study. Other researchers have also reported that elevated procalcitonin is particularly linked to abscess and perforation, further supporting its role as a severity-specific marker, rather than a general diagnostic marker for appendicitis.¹⁵

The current study also supports the use of CRP to assess severity of appendicitis.¹⁶ Although CRP did not perform better than procalcitonin, it did demonstrate some discriminatory power and may still be useful in situations where procalcitonin is not readily available.¹⁷ This is relevant to many public and low-resource hospitals where CRP and WBC count are more widely available and affordable than procalcitonin tests.¹⁸ Therefore, practically, the results indicate that procalcitonin is the best marker, but CRP can be used as an alternative,

particularly when the cost or availability of procalcitonin testing precludes its routine use.¹⁹

The results are also important from a clinical perspective.²⁰ The ability to preoperatively diagnose complicated appendicitis may affect multiple components of care, including the urgency of surgery, the need for more potent antibiotics, the likely surgical complexity, patient and family education, and the level of postoperative care required.¹ A patient with a high procalcitonin and CRP, and clinical signs such as prolonged pain, fever, guarding and rebound tenderness, is more likely to have gangrene, perforation or abscess, and thus may require more urgent surgery and more intensive post-operative management.² On the other hand, patients with lower procalcitonin and CRP levels and less severe clinical symptoms and signs may be more likely to have uncomplicated appendicitis.³ This is especially important as management options for uncomplicated appendicitis broaden, including non-operative management in some cases.⁴ The appendicitis guidelines also highlight that the use of biomarkers should be in combination with clinical scoring and imaging, rather than in isolation.⁵

Limitations of the study

First, it was a single-center study with a small sample of 100 patients, which could affect the applicability of the results to broader patient populations. Second, the study did not include receiver operating characteristic (ROC) curve analysis or biomarker-specific optimized cut-off modeling, which may have improved diagnostic accuracy. Third, only three biomarkers were assessed, and other inflammatory markers such as the neutrophil-to-lymphocyte ratio, immature granulocyte count, platelet indices, or composite clinical scores were not included. Finally, the study evaluated patients with confirmed appendicitis who were treated with surgery, which may not reflect the entire spectrum of patients with equivocal appendicitis who are observed or treated conservatively. However, the study has several strengths. It evaluated three widely available laboratory markers in the same patient population, with operative and histopathological confirmation as the gold standard, included both sexes, and answered a clinically relevant question with important implications for emergency surgical practice¹². Crucially, it is derived from a hospital-based study, which increases its relevance to clinical practice.

This study suggests that serum procalcitonin, in combination with CRP and clinical examinations, may be a useful marker for the early preoperative diagnosis of complicated appendicitis. The use of these markers in emergency surgical practice may help guide risk stratification, timing of surgery and other perioperative decisions. Multicentre studies with larger cohorts are

advised to confirm the findings and to define cut-off levels for the use of these markers.

CONCLUSION

The patients with complicated acute appendicitis had higher levels of serum procalcitonin, CRP and WBC, longer duration of symptoms, and more severe clinical signs than patients with uncomplicated appendicitis. Of the three markers, serum procalcitonin had the highest diagnostic performance, and is therefore the most accurate laboratory marker for distinguishing complicated from uncomplicated acute appendicitis in this population. CRP also demonstrated good diagnostic accuracy, while the WBC count, while useful, was less accurate.

DECLARATION

Conflict of Interest: The authors declare no conflict of interest.

Funding: This research did not receive any external funding.

Author's Contribution: All authors contributed equally in the complication of current study.

Acknowledgments: The authors express their sincere gratitude to all colleagues and participants for their valuable contributions to this study.

REFERENCES

- Cui W, Liu H, Ni H, Qin X, Zhu L. Diagnostic accuracy of procalcitonin for overall and complicated acute appendicitis in children: a meta-analysis. *Ital J Pediatr* 2019;45(1):78.
- Li Y, Zhang Z, Cheang I, Li X. Procalcitonin as an excellent differential marker between uncomplicated and complicated acute appendicitis in adult patients. *Langenbecks Arch Surg* 2020;46(4):853-8.
- Di Saverio S, Podda M, De Simone B, Ceresoli M, Augustin G, Gori A, et al. Diagnosis and treatment of acute appendicitis: 2020 update of the WSES Jerusalem guidelines. *World J Emerg Surg* 2020;15:27.
- Dale L. The use of procalcitonin in the diagnosis of acute appendicitis: a systematic review. *Cureus* 2022;14(10):e30292.
- Dinç T, Sapmaz A, Erkuş Y, Yavuz Z. Complicated or non-complicated appendicitis? That is the question. *Ulus Travma Acil Cerrahi Derg* 2022;28(3):390-94.
- Duyan M, Vural N. Assessment of the diagnostic value of novel biomarkers in adult patients with acute appendicitis: a cross-sectional study. *Cureus* 2022;14(12):e32307.
- Arruzza E, Milanese S, Li LSK, Dizon J. Diagnostic accuracy of computed tomography and ultrasound for the diagnosis of acute appendicitis: a systematic review and meta-analysis. *Radiography (Lond)* 2022;28(4):1127-41.
- Akçiçek M, Ilgar M, Ünlü S. Is acute appendicitis complicated or uncomplicated? Approaching the question via computed tomography. *Acta Radiol* 2023;64(5):1755-64.
- Sikander B, Rosenberg J, Fonnes S. Individual biomarkers in the blood are not yet applicable in diagnosing complicated appendicitis: a scoping review. *Am J Emerg Med* 2023; 67:100-107.
- Gan DEY, Nik Mahmood NRK, Chuah JA, Hayati F. Performance and diagnostic accuracy of scoring systems in adult patients with suspected appendicitis. *Langenbecks Arch Surg* 2023;408(1):267.
- Kaewlai R, Tongchai S, Teerasamit W, Wongsangchan D, Noppakunsomboon N, Khamman P, et al. Validation of scoring systems for the prediction of complicated appendicitis in adults using clinical and computed tomographic findings. *Insights Imaging* 2023;14(1):191.
- Singh A, Pogorelić Z, Agrawal A. Utility of ischemia-modified albumin as a biomarker for acute appendicitis: a systematic review and meta-analysis. *J Clin Med* 2023;12(17):5486.
- Zhang Q, Zhao H, Wang F, Li W, Zhang P. Diagnostic value of laboratory parameters for complicated appendicitis: a two-center study. *Biomed Rep* 2024;20(5):77.
- Saridas A, Vural N, Duyan M, Guven HC, Ertas E, Cander B. Comparison of the ability of newly inflammatory markers to predict complicated appendicitis. *Medicine (Baltimore)* 2024;19(1): 20241002.
- Ha SC, Tsai YH, Koh CC, Hong SG, Chen Y, Yao CL. Blood biomarkers to distinguish complicated and uncomplicated appendicitis in pediatric patients. *J Formos Med Assoc* 2024;123(10):1093-8.
- Düzgün AP, Özel HP, Şahingöz E, Dinç T. Identification of complicated and non-complicated appendicitis: a new Alvarado-based scoring system. *Ulus Travma Acil Cerrahi Derg* 2024;30(2):101-6.
- Şimşek O, Şirolu S, Özkan Irmak Y, Hamid R, Ergun S, Kepil N, et al. Comparative study of imaging features in uncomplicated and complicated acute appendicitis. *Ulus Travma Acil Cerrahi Derg* 2024;30(10):722-8.
- Scheijmans JCG, Bom WJ, Ghori UH, van Geloven AAW, Hannink G, van Rossem CC, et al. Development and Validation of the Scoring System of Appendicitis Severity 2.0. *JAMA Surg* 2024;159(6):642-9.
- Patel H, Kamel M, Cooper E, Bowen C, Jester I. The Variable Definition of "Negative Appendicitis" Remains a Surgical Challenge. *Pediatr Dev Pathol* 2024;27(6):552-8.
- Podda M, Ceresoli M, De Simone B, Fugazzola P, Pata F, Balla A, et al. Diagnosis and Treatment of Acute Appendicitis: 2025 Edition of the World Society of Emergency Surgery Jerusalem Guidelines. *JAMA Surg* 2026; 161(3): 283-95.

Publisher's Note:

Pakistan Journal of Medical & Health Sciences (Pak J Med Health Sci) remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.